



ISTITUTO DI ANALISI DEI SISTEMI ED INFORMATICA
“Antonio Ruberti”
CONSIGLIO NAZIONALE DELLE RICERCHE

D. Naddef, G. Rinaldi

**THE SYMMETRIC
TRAVELING SALESMAN POLYTOPE:
NEW FACETS
FROM THE GRAPHICAL RELAXATION**

R. 609* Marzo 2004

Denis Naddef – LID-IMAG, Institut National Polytechnique de Grenoble - ENSIMAG-Antenne de Montbonnot ZIRST - 51 Avenue Kuntzmann, 38330 Monbonnot St. Martin, France, (denis.naddef@imag.fr).

Giovanni Rinaldi – Istituto di Analisi dei Sistemi ed Informatica “Antonio Ruberti” del CNR, viale Manzoni 30, 00185 Roma, Italy, (rinaldi@iasi.cnr.it).

* Nuova versione del rapporto R. 248, Dicembre 1988

ISSN: 1128–3378

Collana dei Rapporti dell'Istituto di Analisi dei Sistemi ed Informatica "Antonio Ruberti",
CNR

viale Manzoni 30, 00185 ROMA, Italy

tel. ++39-06-77161

fax ++39-06-7716461

email: iasi@iasi.cnr.it

URL: <http://www.iasi.cnr.it>

Abstract

The *path*, the *wheelbarrow*, and the *bicycle* inequalities have been shown in [5] to be facet defining for the *graphical relaxation* of $\text{STSP}(n)$, the polytope of the symmetric traveling salesman problem on an n -node complete graph. We show that these inequalities, and some generalizations of them, define facets also for $\text{STSP}(n)$. In conclusion, we characterize a large family of facet defining inequalities for $\text{STSP}(n)$ that include, as special cases, most of the inequalities currently known to have this property as the *comb*, the *clique tree*, and the *chain* inequalities. Most of the results given here come from a strong relationship of $\text{STSP}(n)$ with its graphical relaxation that we have pointed out in [18], where the basic proof techniques are also described.

Key words: traveling salesman problem, Hamiltonian cycle, polyhedron, facet, linear inequality.

AMS subject classifications: 90C57, 65K05,

1. Introduction

“The most spectacular success of the cutting plane technique has certainly been achieved for the traveling salesman problem” observe Grötschel and Lovász in their survey on Combinatorial Optimization [8]. This success is due mostly to the exploitation of the current (partial) knowledge of the structure of the traveling salesman polytope and is the main motivation for pushing this knowledge a bit further.

In [5], Cornuéjols, Fonlupt, and Naddef define a class of valid inequalities for the symmetric traveling salesman polytope on a graph with n nodes (STSP(n)) known as the *path*, the *wheelbarrow*, and the *bicycle* inequalities (the PWB inequalities for short). They show that these inequalities are facet defining for a relaxation of STSP(n), namely the *graphical relaxation polyhedron* (GTSP(n)).

The main result of this paper is that these inequalities, and many of their generalizations, are facet defining for STSP(n). This result was contained in the research report [16], which had the same title as this article and was widely referenced in the literature, both in research and in survey articles. However, it was very technical and difficult to read, thus was never submitted for publication. This paper is a substantial revision of that report.

The motivation for adding the PWB inequalities to the list of those defining facets of STSP(n), besides the fact that they provide a huge and natural generalization of the *comb* inequalities, comes from the fact that they are useful in a polyhedral cutting plane algorithm for the solution of the traveling salesman problem as shown by Naddef and Thienel in [19, 20] (see also [4]). This experimental evidence confirms the theoretical expectation expressed by Goemans in [6], where he addresses the problem of measuring the quality of a class of inequalities. As a measure, he proposes the ratio of the lower bound for the graphical traveling salesman problem obtained by using all the inequalities of a given class along with the subtour elimination inequalities and the lower bound produced with only the subtour elimination inequalities. The best value, among all the inequalities of GTSP(n) known to date, is obtained for the PWB inequalities.

In [18] it is shown how the two polyhedra STSP(n) and GTSP(n) are very tightly related. On this basis, a technique is developed to prove that some given facet defining inequalities for GTSP(n) are also facet defining for STSP(n). Moreover, sufficient conditions are stated under which the composition of facet defining inequalities introduced in [17] for GTSP(n), also applies to the case of STSP(n). Finally, it is shown how to apply several kinds of liftings to the inequalities that define facets of STSP(n).

In this paper we first use the technique of [18] to prove that the path, the wheelbarrow, and the bicycle inequalities are facet defining for STSP(n) (Section 2). Since these inequalities contain the comb inequalities as a special case, as a byproduct we get another proof that the comb inequalities define facets of STSP(n). Another such proof for comb inequalities can be found in [21], besides, of course, the original proof of Grötschel and Padberg [9, 10].

In [18], an edge cloning operation is defined to extend an inequality to a higher dimensional space, and sufficient conditions under which its application preserves the facet defining property of an inequality are given. Here we apply this operation to the PWB inequalities. The so called *chain inequalities* of [24] are a particular case of the inequalities obtained in this way.

We finally prove that the repeated 2-sum composition of PWB inequalities yields facet defining inequalities. *Clique trees*, with at least one node outside all handles and teeth, are a special case of these inequalities, therefore we get an alternative proof that they are facet defining; the original one was given by Grötschel and Pulleyblank in [11].

4.

Let $G = (V, E)$ be a graph on n nodes. By $e = (u, v)$ we denote the edge of G having u and v as endnodes and we let \mathbb{R}^E be the set of all real vectors whose components are indexed by the edge set E . For every real vector x in \mathbb{R}^E , by x_e , or by $x(u, v)$, we denote the component of x indexed by $e = (u, v)$. For a subset $F \subseteq E$, we let $x(F)$ be the sum $\sum_{e \in F} x_e$. When G is complete, we denote it by $K_n = (V, E_n)$.

With every Hamiltonian cycle H of K_n we associate a unique incidence vector χ^H in \mathbb{R}^{E_n} . The components of χ^H indexed by the edges of H have value 1 while all the other components have value 0. The *symmetric traveling salesman polytope* (also called the *Hamiltonian cycle polytope*) associated with K_n is denoted by $\text{STSP}(n)$ and is the convex hull of the set of the incidence vectors of all the Hamiltonian cycles of K_n .

The description of $\text{STSP}(n)$ with linear inequalities is a classical topic of Polyhedral Combinatorics and has attracted a lot of interest. For the fundamentals of Polyhedral Combinatorics we refer the reader to [22]. While for small values of n ($n \leq 9$) a complete description of the system of inequalities defining facets of $\text{STSP}(n)$ has been generated by means of a computer program (see, e.g., [2]), for arbitrary values of n the knowledge of such a system is far from being complete and it is very unlikely that it will ever be. In the last 40 years many papers appeared in which new valid or facet defining inequalities for $\text{STSP}(n)$ were introduced. We refer to [12], [13], [14], and [15] for a list of them and for further details on the traveling salesman polytope. As new inequalities are discovered, it becomes more and more difficult to keep track of all of them in a unifying framework; moreover the proof techniques are usually specific for each class of inequalities.

This paper aims at giving a compact combinatorial description for a large family of facet defining inequalities that includes most of the known ones as well as at providing a standard proof technique to show that an inequality defines a facet of $\text{STSP}(n)$. As $\text{STSP}(n)$ is not full dimensional, to study its polyhedral structure it is customary to imbed it into a full dimensional polyhedron called a *relaxation*, which is obtained by dropping some conditions on the solution set, and then to find sufficient conditions for an inequality facet defining for the relaxation to maintain such a property for $\text{STSP}(n)$.

The two major relaxations that have been considered in the study of the polyhedral structure of $\text{STSP}(n)$ are the *monotone traveling salesman polytope*, introduced in [7], and the *graphical traveling salesman polyhedron*. Sufficient conditions for a facet defining inequality for the monotone relaxation to be facet defining for $\text{STSP}(n)$ are given in [1].

A desirable property of a relaxation R is that every facet of $\text{STSP}(n)$ be contained in exactly one of the facets of R that do not contain the entire polytope $\text{STSP}(n)$. If this property holds, then there is a one to one correspondence between a subset of the facets of R and all facets of $\text{STSP}(n)$. Unfortunately, the monotone traveling salesman polytope does not have such a property. On the contrary, the graphical traveling salesman polyhedron does. For this reason and for some nice connections with $\text{STSP}(n)$, which will be mentioned in the following, this polyhedron appears to be the most natural and useful relaxation for studying the polyhedral structure of $\text{STSP}(n)$.

We exploit here such connections between $\text{STSP}(n)$ and the graphical traveling salesman polyhedron. The latter polyhedron has been studied in [5, 17], while its connections with $\text{STSP}(n)$ have been studied in [18]. Most of our proofs are based on the results of these three papers.

To formally define the graphical traveling salesman polyhedron, we need the following definitions.

A *multiset of edges* of $G = (V, E)$ is a collection F of elements of E that may contain several copies of the same element. For every element e of E , we call *multiplicity* of e in F the number

of times e appears in F . Clearly, a *set of edges* of G is a multiset where every element has multiplicity 1. Let F_1 and F_2 be two multisets of edges of G and let $F_1 + F_2$ denote the multiset for which the multiplicity of every element is given by the sum of its multiplicities in F_1 and F_2 . By $F + e$ and, if $e \in F$, by $F - e$ we denote the multisets for which the element e has multiplicity one more and one less than in F , respectively. Finally, $t\{e\}$ denotes the multiset containing only the element e with multiplicity t .

Let F be a multiset of edges of $G = (V, E)$. By $G[F]$ we denote the multigraph having node set V and having, for every pair of distinct nodes u and v in V , as many edges with endnodes u and v as the multiplicity of (u, v) in F . For every node v in V the *degree of v in F* is the degree of v in the multigraph $G[F]$, and the *neighbors of v in F* are the neighbors of v in the multigraph $G[F]$. With every multiset F of edges of G we associate a unique representative vector $\chi^F \in \mathbb{R}^E$ by setting χ_e^F equal to the multiplicity of e in F for every $e \in E$. If c is a vector in \mathbb{R}^E , the *c -length of F* , also denoted by $c(F)$, is defined as $c(F) = c\chi^F$. For any two multisets F_1 and F_2 of edges of G , if $\chi^{F_1} \leq \chi^{F_2}$ we say that F_1 is contained in F_2 .

A *spanning closed walk* of a graph $G = (V, E)$ is a multiset W of edges of G such that

- (i) the degree in W of every $v \in V$ is positive and even;
- (ii) $G[W]$ is connected.

We simply use the term *walk* for a spanning closed walk since all our walks are of this kind.

Thus, a Hamiltonian cycle in G is a walk where every node has degree 2, while a walk is not, in general, a Hamiltonian cycle.

Figure 1 shows a graph and a closed walk which is not a Hamiltonian cycle.




Figure 1: Example of a closed walk W in a graph G

The *graphical traveling salesman polyhedron* associated with the graph G , denoted by $\text{GTSP}(G)$ or $\text{GTSP}(n)$ when $G = K_n$, is the convex hull of the set of the representative vectors of all the walks of G and is the polyhedron associated with the *graphical traveling salesman problem*.

If G is connected, then $2T = \{2\{e\} \mid e \in T\}$ is a walk for any spanning tree T of G . Moreover, $W + 2t\{e\}$ is a walk for any walk W , for any edge e of G , and for any nonnegative integer t . Therefore, $\text{GTSP}(G)$ is a full dimensional unbounded polyhedron. Clearly, $\text{GTSP}(n)$ is a relaxation of $\text{STSP}(n)$ (the degree of each node is no longer required to be 2 but only positive and even) and $\text{STSP}(n) = \{x \in \text{GTSP}(n) \mid x(E) = n\}$ (see [5]). Therefore, $\text{STSP}(n)$ is a face of $\text{GTSP}(n)$.

For any inequality $fx \geq f_0$ and for each node $u \in V$, we define the edge set $\Delta_f(u) = \{(v, w) \in E_n \mid u \neq v, u \neq w, f(v, w) = f(u, v) + f(u, w)\}$.

Definition 1.1. An inequality $fx \geq f_0$ defined on \mathbb{R}^{E_n} is said to be tight triangular (abbreviated *TT*) or in tight triangular form (abbreviated *TT form*) if:

- (a) The coefficients f_e satisfy the triangular inequality, i.e., $f(u, v) \leq f(u, w) + f(w, v)$ for every triple u, v, w of distinct nodes in V ;
- (b) $\Delta_f(u) \neq \emptyset$ for all u in V .

In [18] it is proven that, except for the trivial inequalities $x_e \geq 0$ and for the degree inequalities $x(\delta(\{u\})) \geq 2$, all facet defining inequalities of $\text{GTSP}(n)$ are in TT form (here and in the following, by $\delta(U) = \{(u, v) \in E \mid u \in U, v \in V \setminus U\}$ we denote the cocycle of a subset U of V in the graph $G = (V, E)$). Moreover, it is shown there that every non trivial inequality $cx \geq c_0$ facet defining for $\text{STSP}(n)$ has a unique equivalent inequality $fx \geq f_0$ in TT form, up to scaling by a nonnegative constant π , where f and f_0 are defined as follows:

$$\begin{aligned} f(u, v) &= \pi(\lambda_u + \lambda_v + c(u, v)) & \text{for all } (u, v) \in E \\ f_0 &= \pi \left(2 \sum_{u \in V} \lambda_u + c_0 \right), \end{aligned} \tag{1}$$

where $\lambda \in \mathbb{R}^V$ satisfies

$$\lambda_u = \frac{1}{2} \max \{c(v, w) - c(u, v) - c(u, w) \mid u, v, w \in V, u \neq v \neq w\} \quad \text{for all } u \in V. \tag{2}$$

We now show the TT form of the comb inequality, probably the best know inequality of the linear description of $\text{STSP}(n)$. We consider the simplest comb inequality, the one with three teeth. Such an inequality is defined on a subset of vertices H called the *handle* and on three mutually disjoint subsets of vertices T_1, T_2 , and T_3 , called the *teeth*, which intersect H (see Figure 2).




Figure 2: A 3-tooth comb and the coefficients of the inequality in TT form

When first defined by Chvátal [3] and then by Grötschel and Padberg [9] the inequality, where k is odd and stands for the number of teeth (in our example $k = 3$) and where $\gamma(U)$ denotes the edge set $\{(u, w) \mid u, w \in U\}$, was given as

$$x(\gamma(H)) + \sum_{i=1}^k x(\gamma(T_i)) \leq |H| + \sum_{i=1}^k (|T_i| - 1) - (k + 1)/2.$$

After multiplying both sides of the inequality by -1 and applying (2), one obtains for λ_u a value given by half the number of sets (handle and teeth) to which u belongs. Then, by applying (1) (with $\pi = 2$ to produce integral coefficients), one obtains the following version of the inequality in TT form:

$$x(\delta(H)) + \sum_{j=1}^k x(\delta(T_j)) \geq 3k + 1.$$

In the right hand side part of Figure 2 some edges of G are drawn with their coefficients in the TT -form of the comb inequality defined by the sets in the left hand side. The coefficient of any edge e is given by the number of sets whose border is crossed by e . To unclutter the figure, only a few examples of such coefficients are shown.

In [18] it is shown that the TT form of any facet defining inequality for $STSP(n)$ is also facet defining for $GTSP(n)$ ¹. Finally, sufficient conditions are given for an inequality facet defining for $GTSP(n)$ to define a facet of $STSP(n)$.

Based on these results, the process of finding new valid inequalities for $STSP(n)$ and of proving that they are facet defining can go along the following four steps.

1. One first restricts the attention to a (possibly sparse) spanning graph $\bar{G} = (V, \bar{E})$ and proves that an inequality, say $cx \geq c_0$, defines a facet of $GTSP(\bar{G})$. This task may take advantage from the sparsity of \bar{G} and from the fact that using walks, rather than Hamiltonian cycles, simplifies the proofs considerably. Note that \bar{G} only has to be connected, in order to have a full dimensional polyhedron $GTSP(\bar{G})$, and actually may not be Hamiltonian at all. The graph \bar{G} is called the *skeleton* of the inequality (see [17], pp. 373–374).

2. Using a standard sequential lifting procedure, one then extends the inequality $cx \geq c_0$ to become a facet defining inequality for $GTSP(n)$. This is done by choosing an ordering $\langle e_1, e_2, \dots, e_r \rangle$ of the edges of the set $E_n \setminus \bar{E}$ and then, for each $\ell = 1, \dots, r$, by assigning the smallest coefficient to e_ℓ such that the length of the shortest walk in $G_\ell = (V, \bar{E} \cup \bigcup_{t=1}^\ell e_t)$ is c_0 . The resulting inequality is facet defining for $GTSP(n)$ by construction. Different orderings of the edges in $E_n \setminus \bar{E}$ yield, in general, different inequalities. When this is not the case we say that the skeleton is *stable* (see [17]). Examples of a skeleton that is stable and of one that is not are given in Section 2.

3. Once a facet defining inequality for $GTSP(n)$ has been obtained, one tries to show that it defines also a facet of $STSP(n)$ by proving that one of the mentioned sufficient conditions is satisfied.

4. Using the inequalities produced at Step 3 as building blocks, one can apply some operations described in [17] and [18], which produce new inequalities from known ones, to describe in a compact way a large family of facet defining inequalities.

It is easy to see that a TT inequality cannot have negative coefficients. However, it can have coefficients with value zero. A TT inequality $cx \geq c_0$ defined on \mathbb{R}^{E_n} with $c_e > 0$ for all $e \in E_n$ is called *simple*. Simple inequalities have a peculiar geometric property. They define all bounded facets of $GTSP(n)$.

Let $(U : W)$ denote the edge set $\{(u, w) \mid u \in U, w \in W\}$. Suppose we are given a TT inequality $cx \geq c_0$ which is not simple. It is easy to see that V can be partitioned into p sets V^1, \dots, V^p such that:

- (a) $c_e = 0$ for all $e \in \gamma(V^i)$, $i = 1, \dots, p$;
- (b) for all $i \neq j \in \{1, \dots, p\}$, $c_e = c_f > 0$ for $e, f \in (V^i : V^j)$.

Since the edges linking two vertices contained in a same subset V^i have a zero coefficient, these edges do not play any structural role in the definition of the inequality, only the edges linking the various subsets do. In fact, as far as $GTSP(n)$ is concerned, given a facet defining inequality, one can add as many vertices as desired to any subsets V^i and obtain a facet defining inequality for $GTSP(n')$, $n' > n$. This suggests to first study the case in which $|V^i| = 1$ for all i , i.e., to study first simple facet defining inequalities.

The subgraph of G induced by the set obtained by taking one representative node for each of the sets V_i for $i = 1, \dots, p$, is a complete graph K_p on p nodes. The *simple inequality associated with $cx \geq c_0$* is the inequality $\bar{c}\bar{x} \geq c_0$ defined by it on the complete graph K_p with

$$\bar{c}(u_i, u_j) = c_e, \quad e \in (V^i : V^j) \quad \text{for all } 1 \leq i < j \leq p.$$

¹Till very recently, no examples were known of facet defining inequalities for $GTSP(n)$ that are provably not facet defining for $STSP(n)$. In [23] Oswald, Reinelt and Theis exhibit some of such inequalities.

From now on, every time we refer to an inequality of type \mathcal{A} without using the attribute “simple” we mean that the inequality may have zero coefficients and that it is associated to a simple inequality of type \mathcal{A} .

The paper is organized as follows. In Section 2 we show that the simple *path*, *wheelbarrow*, and *bicycle* inequalities (simple PWB for short), proved in [5] to be facet defining for GTSP(n), define also facets of STSP(n).

In Section 3 we extend the result to all PWB inequalities. In Section 4 we prove that the same holds for the a generalization of these inequalities which we call *extended* PWB inequalities. PWB and extended PWB are obtained from the simple PWB inequalities by applying the operations mentioned at the step 4 of our “facet hunting” process described above. In Section 5 we study some compositions of PWB inequalities that yield a large superclass of the clique-tree inequalities.

2. The simple PWB inequalities

The PWB inequalities were first defined in [5]. We give here the alternate definition proposed in [15] that has been more widely used in the recent literature on the traveling salesman problem and, being based on handles and teeth, is more similar to the classical definition of the comb inequalities of which the PWB are a generalization.

Definition 2.1. *A k -PWB configuration is a quadruple $\langle \mathcal{H}, \mathcal{T}, \alpha, \beta \rangle$, where $\mathcal{H} = \{H_r \mid r = 1, \dots, h\}$ and $\mathcal{T} = \{T_i \mid i = 1, \dots, k\}$ are two collections of subsets of V called the handles and the teeth of the configuration, respectively, and α and β are integer vectors of h and k components, respectively. Each handle H_r has associated the component α_r and each tooth T_i has associated the component β_i . The vectors α and β satisfy the Equality Interval Property (see Definition 2.2 below), moreover the handles and the teeth satisfy the following conditions:*

$$H_r \subset H_{r+1} \quad \text{for } 1 \leq r \leq h-1 \quad (3)$$

$$H_1 \cap T_i \neq \emptyset \quad \text{for } i = 1, \dots, k \quad (4)$$

$$T_i \setminus H_h \neq \emptyset \quad \text{for } i = 1, \dots, k \quad (5)$$

$$T_i \cap T_j = \emptyset \quad \text{for } 1 \leq i < j \leq k \quad (6)$$

$$(H_{r+1} \setminus H_r) \setminus \bigcup_{i=1}^k T_i = \emptyset \quad \text{for } 1 \leq r \leq h-1 \quad (7)$$

The above conditions state that the handles are linearly nested (3), that the innermost handle intersects all teeth (4), that each tooth has at least one node not contained in any handle (5), that the teeth are pairwise disjoint (6), and finally that the node of each handle is contained either in H_1 or in a tooth (7).

In Figure 3 an example of a PWB configuration with three teeth and six handles is shown. The black filled points in the figure represent nonempty sets of nodes while the white filled points represent possibly empty sets of nodes. The union of all these node sets gives V . It is easy to verify that the handles and the teeth satisfy the conditions (3)-(7). In particular, by Condition (7), the nodes that are not contained in any tooth can either belong to H_1 (in the figure the set of these nodes is marked as Y) or belong to the complement of H_h (the set of these nodes is marked as Z).

To complete Definition 2.1, we have to define the *Equality Interval Property*. To this purpose, we need the notion of *handle interval* relatively to a given tooth T_i . A *handle interval* relatively to a given tooth T_i of a k -PWB configuration is the index set $R \subseteq \{1, \dots, h\}$ of a maximal set




Figure 3: Example of a PWB configuration

of handles which have the same intersection with T_i . By definition, a handle interval is made of consecutive indices. Moreover, the index set $\{1, \dots, h\}$ is partitioned into s_i handle intervals relatively to a tooth T_i . For example, in the PWB configuration of Figure 3, the handle intervals relatively to T_1 are $\{1, 2, 3\}$ and $\{4, 5, 6\}$, those relatively to T_3 are $\{1\}$, $\{2, 3\}$, $\{4, 5\}$, and $\{6\}$.

Definition 2.2. *The vectors α and β of a k -PWB configuration $\langle \mathcal{H}, \mathcal{T}, \alpha, \beta \rangle$ satisfy the Equality Interval Property if $\sum_{r \in R} \alpha_r = \beta_i$ holds for each tooth T_i and for each handle interval R relative to T_i .*

It is easy to check that the integers associated with each handle and tooth in Figure 3 define vectors α and β that do satisfy the Equality Interval Property, thus the one shown is indeed a PWB configuration.

Definition 2.3. *A k -PWB inequality associated with the k -PWB configuration $\langle \mathcal{H}, \mathcal{T}, \alpha, \beta \rangle$ is the inequality*

$$\sum_{r=1}^h \alpha_r x(\delta(H_r)) + \sum_{i=1}^k \beta_i x(\delta(T_i)) \geq (k+1) \sum_{r=1}^h \alpha_r + 2 \sum_{i=1}^k \beta_i \quad (8)$$

Note that the coefficient of any edge e is easily computed by adding up the coefficient of all the sets whose border is crossed by e .

The coefficient vector of inequality (8) is the positive weighted sum of incident vectors of cocycles in G . Thus, each coefficient satisfies the triangular inequality. It is not difficult to verify that also condition (b) of Definition 1.1 is satisfied, hence (8) is in TT form.

A comb inequality is a special case of a PWB-inequality. Its configuration has $|\mathcal{H}| = 1$ and the components of α and β have all value 1.

Remark 2.4. *If the number of teeth k is even, then the inequality (8) is not valid for the $STSP(n)$.*

Definition 2.5. A k -PWB configuration (inequality) is called a k -path, a k -wheelbarrow, or a k -bicycle configuration (inequality) depending on whether both the subsets $H_1 \setminus \bigcup_{i=1}^k T_i$ and $(V \setminus H_1) \setminus \bigcup_{i=1}^k T_i$ are nonempty, or only one of two is empty, or they are both empty.

The Equality Handle Interval property is very binding, and for an arbitrarily chosen quadruple $\langle \mathcal{H}, \mathcal{T}, \alpha, \beta \rangle$ that satisfies the conditions (3)-(7) there are very little chances that it is satisfied. For generating all possible k -PWB configuration, a different approach might be more useful where the elements of the quadruple $\langle \mathcal{H}, \alpha, \mathcal{T}, \beta \rangle$ are generated by the following simple procedure.

PROCEDURE 2.6. Input: any odd $k \geq 3$, any k -tuple of positive integers (n_1, \dots, n_k) , with $n_i \geq 2$ for $i \in \{1, \dots, k\}$ and any partition of V into the sets $S_Y, S_Z, S_1^1, \dots, S_{n_1}^1, S_1^2, \dots, S_{n_2}^2, \dots, S_1^k, \dots, S_{n_k}^k$, where only S_Y and S_Z are allowed to be empty. Output: a quadruple $\langle \mathcal{H}, \mathcal{T}, \alpha, \beta \rangle$ satisfying the Equality Handle Interval property and (3)-(7).

1. let ζ be the least common multiple of the integers $n_1 - 1, \dots, n_k - 1$;
2. for $i = 1, \dots, k$, let $T_i = \bigcup \{S_j^i \mid j = 1, \dots, n_i\}$ and $\beta_j = \zeta / (n_i - 1)$;
3. let $H_1 = \bigcup \{S_1^i \mid i = 1, \dots, k\} \cup \{Y\}$ and $\alpha_1 = \min\{\beta_i \mid i = 1, \dots, k\}$;
4. for $i = 1, \dots, k$, let $j_i = 1$ and $\beta'_i = \beta_i - \alpha_1$;
5. let $\ell = 1$;
6. while $j_i < n_i - 1$ for some $i \in \{1, \dots, k\}$ do the following:
 - (a) increment ℓ by 1;
 - (b) for any i such that $\beta'_i = 0$, let $\beta'_i = \beta_i$ and increment j_i by 1;
 - (c) let $H_\ell = H_{\ell-1} \bigcup \{S_{j_i}^i \mid i = 1, \dots, k\}$ and $\alpha_\ell = \min\{\beta'_i \mid i = 1, \dots, k\}$;
 - (d) for $i = 1, \dots, k$, decrement β'_i by α_ℓ ;

In the case of Figure 3, let us set $k = 3$ and consider the partition of V given by the sets represented by the black and white filled points. Let S_Y and S_Z be the sets marked as Y and Z , respectively. Finally, for $j = 1, \dots, k$, let $S_1^j, \dots, S_{n_j}^j$ be the sets contained in the tooth T_j and ordered from the top to the bottom. It is easy to verify that the above procedure generates the teeth, the handles and the coefficients α and β as those shown in the drawing.

If the endnodes of an edge of G belong to the same handles and teeth of a k -PWB configuration, then the coefficient of this edge is zero in (8). On the contrary, if there is a handle or a tooth that contains only one of the endnodes, then the edge belongs to one of the cocycles of the handles and of the teeth and its coefficient is positive in (8). Consequently, if there are no two nodes of G that are contained in the same handles and teeth of a k -PWB configuration, the corresponding k -PWB inequality is simple. We call *simple* the k -PWB configuration in this case. For example, the PWB configuration of Figure 3 is simple if each black filled point represents a single node of V and each white filled point represents a single node of V or the empty set. In this section we only consider simple k -PWB configurations and inequalities.

A simple k -PWB configuration induces a labeling of the nodes of G that will be widely used throughout the paper. If $H_1 \setminus \bigcup_{i=1}^k T_i$ is nonempty, its only node is labeled Y ; similarly, if $(V \setminus H_1) \setminus \bigcup_{i=1}^k T_i$ is nonempty, its only node is labeled Z . The n_i nodes of a tooth T_i can be linearly ordered in such a way that $w \in T_i$ precedes $v \in T_i$ if it is contained in more handles

than v . Each node of T_i is labeled u_j^i , where j is the position of the node in such an ordering. Therefore, u_1^i is the node contained in H_1 and $u_{n_i}^i$ is contained in no handles. The value of the superscript of a label u_j^i is defined to be $((i-1) \bmod k) + 1$, thus, for example, $u_j^{k+i} = u_j^i$. Note that this labeling is in agreement with the names given to the sets S_j^i in Procedure 2.6. Consequently, given any odd $k \geq 3$ and any k -tuple of positive integers (n_1, \dots, n_k) , with $n_i \geq 2$ for $i \in \{1, \dots, k\}$ and the labeling

$$\{Y, Z\} \cup \{u_j^i \mid j \in \{1, \dots, n_i\}, i \in \{1, \dots, k\}\}, \quad (9)$$

Procedure 2.6 uniquely identifies a k -PWB configuration $\langle \mathcal{H}, \mathcal{T}, \alpha, \beta \rangle$. Consequently, we refer to the labeling (9) as to a k -PWB configuration.

Depending on whether a k -PWB is a k -path (labels Y and Z are present), a k -wheelbarrow (only label Z is present), or a k -bicycle (Y and Z are missing) configuration, the labeling (9) is denoted by $P(n_1, \dots, n_k)$, $W(n_1, \dots, n_k)$, or $B(n_1, \dots, n_k)$, respectively.

For convenience we label $u_0^i = Y$, and $u_{n_i+1}^i = Z$, for all $i = 1, \dots, k$. The nodes Y and Z are called the *odd nodes* of the configuration, all the other nodes are called *even* (see Figure 4).




Figure 4: The labeling induced by the 3-path configuration of Figure 3 and the skeleton of the corresponding 3-path inequality

In the following, to simplify the notation, by π^i we denote the edge set

$$\pi^i = \{(u_j^i, u_{j+1}^i) \mid j \in \{1, \dots, n_i - 1\}\} \quad \text{for } i \in \{1, \dots, k\}.$$

An edge of any set π^i or any edge (u_0^i, u_1^i) or $(u_{n_i}^i, u_{n_i+1}^i)$ for $i \in \{1, \dots, k\}$ is called a *path edge*. By C_Y and C_Z we denote the cycles whose edge sets are $\{(u_1^i, u_1^{i+1}) \mid i \in \{1, \dots, k\}\}$ and $\{(u_{n_i}^i, u_{n_i+1}^{i+1}) \mid i \in \{1, \dots, k\}\}$, respectively. An edge of any of these two cycles is called a *cycle edge*. An edge of the type (u_1^i, u_1^r) or $(u_{n_i}^i, u_{n_i}^r)$ with $2 \leq ((r-i) \bmod k) \leq k-2$ is called a *chord* (a *short chord* if one of the two inequalities holds with equality) of the cycle C_Y or C_Z , respectively. Recall that paths (teeth) indices are taken modulo k .

Now it is easy to relate the definition above with the following original definition of the path inequality given in [5]:

Definition 2.7. *The simple k -path inequality associated with $P(n_1, \dots, n_k)$ is the following inequality on \mathbb{R}^{E_n} , with $n = 2 + \sum_{i=1}^k n_i$, in which $\eta_i = \frac{1}{n_i-1}$ for $i \in \{1, \dots, k\}$:*

$$cx \geq c_0 = k + 1 + 2 \sum_{i=1}^k \eta_i, \quad (10)$$

where

$$c_e = \begin{cases} |j - q|\eta_i & \text{for } e = (u_j^i, u_q^i), j \neq q, \\ & i \in \{1, \dots, k\}, \\ & j, q \in \{0, \dots, n_i + 1\}, \\ \eta_i + \eta_r + |(j - 1)\eta_i - (q - 1)\eta_r| & \text{for } e = (u_j^i, u_q^r), i \neq r, \\ & i, r \in \{1, \dots, k\}, \\ & j \in \{1, \dots, n_i\}, \\ & q \in \{1, \dots, n_r\}, \\ 1 & \text{for } e = (Y, Z) \end{cases} \quad (11)$$

The coefficient vectors of the simple k -wheelbarrow and of the simple k -bicycle inequalities are restrictions of the coefficient vector of a simple k -path inequality, obtained by removing all the edges incident with Y and all the edges incident with Y and Z , respectively.

The coefficients of the equation (8) are exactly those of Definition 2.7 multiplied by the least common multiple of the numbers $n_i - 1$, $i = 1, \dots, k$. Therefore, the two inequalities of the definitions 2.5 and 2.7 define the same face of $\text{STSP}(n)$.

If $n_i = p$ for $i \in \{1, \dots, k\}$, the simple path, wheelbarrow, and bicycle configurations and their corresponding inequalities are called p -regular. The 2-regular PWB inequalities are the comb inequalities.

For the special case of regular configurations, the coefficients of the associated inequalities can be written as follows:

$$c_0 = (k + 1)(p + 1) - 2, \quad (12)$$

$$c_e = \begin{cases} |j - q| & \text{for } e = (u_j^i, u_q^i), i \in \{1, \dots, k\}, \\ & j \neq q, j, q \in \{0, \dots, p + 1\}, \\ |j - q| + 2 & \text{for } e = (u_j^i, u_q^r), i \neq r, i, r \in \{1, \dots, k\}, \\ & j, q \in \{1, \dots, p\}, \\ p - 1 & \text{for } e = (Y, Z). \end{cases} \quad (13)$$

2.1. The GTSP case

In [5] it is shown that the k -path, the k -wheelbarrow and the k -bicycle inequalities define facets of $\text{GTSP}(G)$, provided that G has a stable skeleton of the inequality as a subgraph. The stable skeleton for a k -path is given by the union of all the path edges (see an example in Figure 4), for the k -wheelbarrow it is given by the union of all the path edges and C_Y , and for the k -bicycle it is given by the union of all the path edges, C_Y , C_Z , and the short chords of these two cycles.

We give here a proof that the k -bicycle inequalities define facets of $\text{GTSP}(n)$ which is slightly different from that of [5] as it also specifies a special set of tight walks that is used in the following proofs.

We make use the following result proven in [5] (Theorem 3.4):

Theorem 2.8. *The simple bicycle inequalities are valid for $\text{GTSP}(n)$.*

To prove the next theorem we need a few more definitions and a simple lemma.

A walk W is said to be *minimal* if it does not contain another walk. For every inequality $fx \geq f_0$ in \mathbb{R}^{E_n} , valid for $\text{STSP}(n)$ or for $\text{GTSP}(G)$, we call *tight* the Hamiltonian cycles and the minimal walks whose representative vectors satisfy the inequality with equality. The set of tight Hamiltonian cycles and tight walks for f are denoted by \mathcal{H}_f^- and \mathcal{W}_f^- , respectively. The

dimension of $\text{GTSP}(G)$ is $|E|$ (i.e., $\text{GTSP}(G)$ is full dimensional) as long as G is connected. A *walk basis* of an inequality $cx \geq c_0$ defining a facet of $\text{GTSP}(G)$ is any set \mathcal{B}_c of $|E|$ walks in \mathcal{W}_c^- whose representative vectors are linearly independent. Note that linear and affine independence of the representative vectors are equivalent since the zero vector does not belong to $\text{GTSP}(G)$. A *Hamiltonian cycle basis* of an inequality $cx \geq c_0$ defining a facet of $\text{STSP}(n)$ is a set \mathcal{C}_c of $|E_n| - |V|$ Hamiltonian cycles in \mathcal{H}_c^- whose representative vectors are linearly independent.

If z is a vector in \mathbb{R}^{E_n} , by $z_{\bar{E}}$ we denote its restriction (projection) to the subspace $\mathbb{R}^{\bar{E}}$ for $\bar{E} \subset E_n$.

Lemma 2.9. *Let $G = (V, \bar{E})$ be a graph with $\bar{E} \subset E_n$, let e^* be an edge in $E_n \setminus \bar{E}$ and $fx \geq f_0$ be a valid inequality for $\text{GTSP}(n)$. If $f_{\bar{E}}x_{\bar{E}} \geq f_0$ is facet defining for $\text{GTSP}(G)$ and there exists a walk of $\mathcal{W}_{\bar{f}}^-$ containing e^* and only edges of \bar{E} , then the inequality $f_{E'}x_{E'} \geq f_0$ is facet defining for $\text{GTSP}(G')$, where $E' = \bar{E} \cup \{e^*\}$ and $G' = (V, E')$.*

Proof. Let \mathcal{B} be a walk basis of $f_{\bar{E}}x_{\bar{E}} \geq f_0$ and let W be a walk of $\mathcal{W}_{\bar{f}}^-$ containing e^* and only edges of \bar{E} . Then $\mathcal{B} \cup \{W\}$ is a set of walks whose representative vectors are linearly independent, i.e., it is a walk basis for $f_{E'}x_{E'} \geq f_0$. \square

The use of Lemma 2.9 is quite evident in our context. Given the stable skeleton \bar{G} , once we prove that the inequality is facet defining for $\text{GTSP}(\bar{G})$, it is straightforward to prove that it is facet defining for $\text{GTSP}(n)$.

In the following theorem and quite often throughout the paper we make use of some special walks given by the following

Definition 2.10. *For $i \in \{1, \dots, k\}$ and $j \in \{1, \dots, n_i - 1\}$, we define (see Figure 5)*

$$\begin{aligned} W_A^i &= 2\pi^i \cup \left\{ (u_{n_s}^s, u_{n_{s+1}}^{s+1}) \mid s \in \{i, i+2, \dots, i+k-1\} \right\} \\ &\quad \cup \left\{ (u_1^s, u_1^{s+1}) \mid s \in \{i+1, i+3, \dots, i+k-2\} \right\} \\ &\quad \cup \left\{ \pi^\ell \mid \ell \in \{1, \dots, k\}, \ell \neq i \right\}, \end{aligned}$$

$$\begin{aligned} W_B^i &= 2\pi^i \cup \left\{ (u_1^s, u_1^{s+1}) \mid s \in \{i, i+2, \dots, i+k-1\} \right\} \\ &\quad \cup \left\{ (u_{n_s}^s, u_{n_{s+1}}^{s+1}) \mid s \in \{i+1, i+3, \dots, i+k-2\} \right\} \\ &\quad \cup \left\{ \pi^\ell \mid \ell \in \{1, \dots, k\}, \ell \neq i \right\}, \end{aligned}$$

$$W_C^i(j) = W_A^i - (u_1^{i-2}, u_1^{i-1}) - 2(u_j^i, u_{j+1}^i) + (u_1^{i-2}, u_1^i) + (u_1^{i-1}, u_1^i),$$

$$W_D^i(j) = W_A^i - (u_1^{i+1}, u_1^{i+2}) - 2(u_j^i, u_{j+1}^i) + (u_1^i, u_1^{i+2}) + (u_1^i, u_1^{i+1}),$$

$$W_E^i = W_B^i - (u_1^i, u_1^{i+1}) - (u_1^{i+2}, u_1^{i+3}) + (u_1^i, u_1^{i+2}) + (u_1^{i+1}, u_1^{i+3}).$$

The walks W_A^i , W_B^i , $W_C^i(2)$, $W_D^i(2)$, and W_E^i are shown in Figures 5(a), 5(b), 5(c), 5(d), and 5(e), respectively, for both the case $k = 3$ and $k \geq 5$. Observe that all the edges of these walks are either path edges, or cycle edges, or short chords of the cycle C_Y . To unclutter the figures, all the edge sets π^i have cardinality 1 in most of them. However, any path edge may be replaced by a path with an arbitrary number of edges. The nodes u_1^i for $i \in \{1, \dots, k\}$ are those at the top of the drawings. Moreover, we have operated in the figures a circular permutation of the paths (teeth). For example, in the figures 5(a) and 5(b) the teeth are ordered from left to right




Figure 5: The tight walks of Definition 2.10

with the indices $i - 1, i, \dots, k, 1, \dots, i - 2$. In the figures 5(d) and 5(e) the order is $i, i + 1, \dots, k, 1, \dots, i - 1$. To extend the walks shown in the figures to the case when k is greater than 5, split a cycle edge into three edges, then remove the central edge (marked with a cross in the figures), and finally connect its endnodes by the path marked by dotted lines in the figures. In the following we will use the same convention for all the figures involving PWB configurations.

Remark 2.11. *The walks of Definition 2.10 satisfy a bicycle inequality at equality. For the sake of simplicity, in the following we refer to these walks also dealing with wheelbarrow and path inequalities. In these cases, though, we will always assume that each of these walks be modified by inserting each required odd node, by removing any cycle edge (of the cycle associated to the odd node) from the walk, and by adding an edge from each of its endnodes to the odd node.*

Theorem 2.12. *A simple bicycle inequality $cx \geq c_0$ associated with the simple k -bicycle configuration $B(n_1, \dots, n_k)$ is facet defining for $GTSP(n)$ and has a walk basis \mathcal{B}_c with the following properties:*

- (a) *Every walk of \mathcal{B}_c intersects the edge set of the cycle C_Y .*
- (b) *Every walk of \mathcal{B}_c intersects the edge set of the cycle C_Z .*

- (c) For each edge $e \in C_Y$ there exists $e' \in C_Y$ and a walk $W \in \mathcal{B}_c$ containing both e and e' .
(d) For each edge $e \in C_Z$ there exists $e' \in C_Z$ and a walk $W \in \mathcal{B}_c$ containing both e and e' .

Proof. We prove first that the inequality $c_S x_S \geq c_0$ is facet defining for $\text{GTSP}(G)$, where $G = (V, S)$ is the subgraph of K_n whose edges are all the path edges, all the cycle edges, and all the short chords of the cycle C_Y .

The inequality $c_S x_S \geq c_0$ is valid by Theorem 2.8 and is supporting since all the walks of Definition 2.10 belong to \mathcal{W}_c^- . We show that the set of all these walks constitutes a basis \mathcal{B}_1 for $c_S x_S \geq c_0$.

Let $f x \geq c_0$ be an inequality defining a facet of $\text{GTSP}(G)$ that contains the face defined by $c_S x_S \geq c_0$, i.e., such that $\mathcal{W}_c^- \subseteq \mathcal{W}_f^-$.

From $f(W_C^i(j')) = f(W_C^i(j''))$ for $j' \neq j'' \in \{1, \dots, n_i - 1\}$ it follows that

$$f(u_{j'}^i, u_{j'+1}^i) = f(u_{j''}^i, u_{j''+1}^i) = g_i \quad \text{for } i \in \{1, \dots, k\}, \quad (14)$$

$$j' \neq j'' \in \{1, \dots, n_i - 1\}.$$

For notational convenience we set $b_i = f(u_{n_i}^i, u_{n_i+1}^i)$, $d_i = f(u_1^i, u_1^{i+1})$, $e_i = f(u_1^i, u_1^{i+2})$, for




Figure 6: The coefficients of some relevant edges

$i \in \{1, \dots, k\}$ (see Figure 6). From $f(W_A^i) = f(W_B^{i+1})$ for $i \in \{1, \dots, k\}$, it follows that

$$(n_i - 1)g_i + b_i = (n_{i+1} - 1)g_{i+1} + d_i \quad \text{for } i \in \{1, \dots, k\}. \quad (15)$$

From $f(W_B^i) = f(W_A^{i+1})$ for $i \in \{1, \dots, k - 1\}$, it follows that

$$(n_i - 1)g_i + d_i = (n_{i+1} - 1)g_{i+1} + b_i \quad \text{for } i \in \{1, \dots, k - 1\}. \quad (16)$$

The equations (15) and (16) imply

$$b_i = d_i \quad \text{for } i \in \{1, \dots, k - 1\} \quad (17)$$

and

$$g_i = \frac{h}{n_i - 1} \quad \text{for } i \in \{1, \dots, k - 1\}. \quad (18)$$

From $f(W_A^{i+1}) = f(W_C^{i+1})$ for $i \in \{1, \dots, k\}$ it follows that

$$e_{i-1} = d_{i-1} - d_i + \frac{2h}{n_{i+1} - 1} \quad \text{for } i \in \{1, \dots, k\}, \quad (19)$$

and from $f(W_C^{i+1}) = f(W_D^{i-1})$ for $i \in \{2, \dots, k\}$ we have

$$d_i = d_{i-1} + \frac{h}{n_i + 1} - \frac{h}{n_i - 1} \quad \text{for } i \in \{2, \dots, k\}. \quad (20)$$

If $k = 3$ then $d_1 = e_2$, otherwise from $f(W_E^1) = f(W_B^1)$ it follows that $d_1 + d_3 = e_1 + e_2$ and from (19) we have

$$d_i = \frac{h}{n_i - 1} + \frac{h}{n_{i+1} - 1} \quad \text{for } i \in \{1, \dots, k\}. \quad (21)$$

If for the bicycle inequality $cx \geq c_0$ we use the definitions 2.1 and 2.3, with the vectors α and β computed with Procedure 2.6, then setting $h = r\zeta$, where ζ is as defined in Procedure 2.6, one can see that $f_e = rc_e$, for e in the skeleton. Since we chose the same right hand sides for both the inequalities, we have $r = 1$. Similarly, if we use Definition 2.7, then setting $r = 1/\zeta$, i.e., setting $h = 1$, we obtain the same result. Therefore, $f \equiv c_S$, the inequality $c_S x_S \geq c_0$ is facet defining for $\text{GTSP}(G)$, and the $|S|$ walks of the set \mathcal{B}_1 are linearly independent.




Figure 7: Tight walks containing an edge across two paths and only edges of a stable skeleton

To complete the proof, we take a sequence $\langle e_1, e_2, \dots, e_s \rangle$ of the edges in $E_n \setminus S$ and we construct the sequences of nested edge sets F_0, F_1, \dots, F_s , with $F_0 = S$ and $F_\ell = F_{\ell-1} \cup \{e_\ell\}$, for $\ell = 1, 2, \dots, s$. Then for every e_ℓ we exhibit a walk of \mathcal{W}_c^- containing e_ℓ and only edges in $F_{\ell-1}$. Consequently, by Lemma 2.9, $cx \geq c_0$ defines a facet of $\text{GTSP}(n)$. We describe now these walks.

Let $e = (u_j^i, u_q^r)$ be any edge of $E_n \setminus S$ and let $d(e, k) = \min\{(i - r) \bmod k, (r - i) \bmod k\}$. Depending on the value of $d(e, k)$ we have different kinds of walks. Clearly, $0 \leq d(e, k) \leq (k - 1)/2$.

For an edge $e = (u_j^i, u_{j+q}^i)$ the value of $d(e, k)$ is zero and we consider the walk

$$W_0(e) = W_A^i \setminus \{(u_{j+t}^i, u_{j+t+1}^i) \mid t = 1, \dots, q-1\} + e.$$

All the edges of $W_0(e)$, except e , belong to S .

For $d(e, k) = 1, \dots, 4$ the corresponding walks $W_1(e), \dots, W_4(e)$ are shown in Figure 7(a), \dots , (d), respectively. The reader should not be confused by the fact that in the figures the endnodes of $e = (u_j^i, u_q^r)$ always have $1 < j < n_i$ and $1 < q < n_r$. Actually, the walks shown can be easily modified to accommodate the cases for all possible values of j and q in the sets $\{1, \dots, n_i\}$ and $\{1, \dots, n_r\}$, respectively, as long as the inequality $(q-1)\eta_r \geq (j-1)\eta_i$ is satisfied. If this is not the case, one can consider the mirror images of the walks of Figure 7, which would correspond to take the opposite orientation of the cycles C_Y and C_Z to draw the walks.

The walks corresponding to a chord of the cycle C_Y (when $j = q = 1$) and to a chord of the cycle C_Z (when $j = n_i$ and $q = n_r$) are of the kind of those shown in Figure 7. All walks $W_\ell(e)$, for $\ell = 0, \dots, 4$, have the edge e and only edges in S .

If $k \leq 9$, then $d(e, k)$ is never greater than 4. Therefore, the above walks are sufficient to complete the proof that the inequality is facet defining, no matter how the sequence of the edges in the set $E_n \setminus S$ is chosen.

It is easy to verify that if $d(e, k) \geq 5$ it is not possible to construct a walk that uses only e and edges in S . However it is always possible to construct a walk $W_5(e)$ that contains e , some edges in S and short chords of the cycle C_Z . Such a walk is shown in Figure 8, where k has any value greater than 11 and $d(e, k)$ has any value greater than 5. To apply Lemma 2.9, the edges of $E_n \setminus S$ have to be ordered in such a way that any edge e with $d(e, k) \geq 5$ follows all the short chords of cycle C_Z in the sequence $\langle e_1, e_2, \dots, e_s \rangle$.




Figure 8: Tight walk made of an edge across two paths, edges of a stable skeleton and short chords of C_Z

The union of \mathcal{B}_1 and all the walks associated with the edges e_1, e_2, \dots, e_s is a walk basis \mathcal{B}_c of the inequality $cx \geq c_0$. It is easy to verify, by inspection, that all its walks intersect both the cycles C_Y and C_Z .

Finally, for any edge $e = (u_1^i, u_1^{i+1})$ of the cycle C_Y , the walk W_B^i contains e and the edge (u_1^{i-1}, u_1^i) , that also belongs to C_Y . Analogously, for any edge $e = (u_{n_i}^i, u_{n_i+1}^{i+1})$ of the cycle C_Z , the walk W_A^i contains e and the edge $(u_{n_i-1}^{i-1}, u_{n_i}^i)$, that also belongs to C_Z . \square

Remark 2.13. As a byproduct of Theorem 2.12, we have that a simple bicycle inequality is facet defining for $GTSP(G)$, where $G = (V, S)$ is the subgraph of K_n whose edges are all the path edges, all the cycle edges, and all the short chords of the cycle C_Y . This result is a bit stronger than that given in [5], where G is required to have also all the short chords of the cycle C_Z .

As it is claimed in the proof of Theorem 2.12, the skeleton (V, S) (where S is the union of the path edges, the edge cycles, and the short chords of C_Y) is stable only for $k \leq 9$. One may wonder what happens for $k \geq 11$, or, more precisely, what happens if the lifting sequence described in the proof of the theorem starts with an edge e with $d(e, k) \geq 5$. If the edge e is a chord of C_Y , then it is still possible to find a tight walk of the bicycle inequality made of e and of edges in S (see Figure 9(a)). This implies that the lifting coefficient resulting for e is the




Figure 9: Tight walks containing a long chord and only edges of a stable skeleton

same as given in Definition 2.7. The same is true if e is a chord of C_Z , but only for $k = 11$ (see Figure 9(b)). However, this is no longer true in general; e.g., take the case where $n_i = 2$ for all i and $k = 11$. If one starts the lifting sequence with $e = (u_1^1, u_2^6)$, then the result is $c_e = 1$; if (u_1^{11}, u_2^6) is the second edge in the sequence, the coefficient is again 1. At this point, no matter how the remaining edges are ordered, the coefficients they get either are as in Definition 2.7, or have value 4. The edges having a coefficient that differs from the one of Definition 2.7 are shown in Figure 10, where the edges with coefficient equal to 1 and 4 are drawn with dashed and solid lines, respectively (note that, to unclutter Figure 10, the arrangement of the nodes has been changed with respect to Figure 9). Thus, the inequality in this way described by Figure 10 and Definition 2.7 is a new facet defining inequality for STSP(22) that shares the same skeleton (V, S) with the bicycle inequality. On the other hand, $(V, S \cup \{(u_1^1, u_2^6), (u_1^{11}, u_2^6)\})$ is a stable skeleton for the new inequality.

2.2. The STSP case

For $u \in V$, we say that a walk W is *almost Hamiltonian in u* if u has degree 4 in W and every other node of V has degree 2 in W .

A basis \mathcal{C}_c of an inequality $cx \geq c_0$, defining a facet of GTSP(n), is called *canonical* if it contains $|E_n| - n$ Hamiltonian cycles and n almost Hamiltonian walks (i.e., one for each $u \in V$).

The following notion of a set of edges being c -connected in a given node, is the key for the sufficient conditions, given in [18], for an inequality facet defining for GTSP(n) to preserve such a property for STSP(n).

Let $e = (u, v)$ and $f = (w, y)$ be two distinct edges in E_n . We say that e and f are c -adjacent if they belong to a tight Hamiltonian cycle $H \in \mathcal{H}_c^-$. Let z be a node in V ; we say that e and f




Figure 10: Edges in which a new facet defining inequality differs from an 11-bicycle

are c -adjacent in z if:

- (i) e and f belong to $\Delta_c(z)$;
- (ii) there exists a walk $W_z \in \mathcal{W}_c^-$ almost Hamiltonian in z that contains the edges (z, u) , (z, v) , (z, w) , and (z, y) ;
- (iii) $W_z - (z, u) - (z, v) + e$ is a Hamiltonian cycle (and such is also $W_z - (z, w) - (z, y) + f$).

A set of edges $J \subseteq E_n$ is said to be c -connected if for every pair of distinct edges f_1 and $f_2 \in J$ there exists a sequence of t edges e_1, \dots, e_t in J , with $e_1 \equiv f_1$ and $e_t \equiv f_2$, where e_i is c -adjacent to e_{i+1} , for $i = 1, \dots, t-1$. A set of edges $J \subseteq E_n$ is said to be c -connected in z if for every pair of distinct edges f_1 and $f_2 \in J$ there exists a sequence of t edges e_1, \dots, e_t in E_n (not necessarily belonging to J), with $e_1 \equiv f_1$ and $e_t \equiv f_2$, where e_i and e_{i+1} are c -adjacent in z , for $i = 1, \dots, t-1$. Observe that the notion of c -connectedness in z is “weaker” than the one of c -connectedness, in the sense that, contrarily to what happens for the usual concept of connectivity, in this case every subset of a set c -connected in z is c -connected in z as well.

Lemma 2.14 (Lemma 2.14 [18]) *Let $cx \geq c_0$ be a TT inequality defining a facet of $GTSP(n)$. If $\Delta_c(u)$ is c -connected in u for every $u \in V$, then $cx \geq c_0$ has a canonical basis, thus it is facet-defining for $STSP(n)$.*

For every ordered triple $\langle u, v, w \rangle$ of distinct nodes in V , we call *shortcut* on $\langle u, v, w \rangle$ the vector $s_{uvw} \in \mathbb{R}^{E_n}$ defined by

$$s_{uvw}(e) = \begin{cases} 1 & \text{if } e = (v, w) \\ -1 & \text{if } e \in \{(u, v), (u, w)\} \\ 0 & \text{otherwise.} \end{cases}$$

Adding a shortcut to a walk that is tight for an inequality can produce a new tight walk with fewer edges. More precisely we have the following

Lemma 2.15 (Lemma 2.5 [18]) *Let $cx \geq c_0$ be a tight triangular inequality which is supporting for $GTSP(n)$, and let $W \in \mathcal{W}_c^-$ be a walk having $t > n$ edges and containing the edge e . For every node $u \in V$ with degree $k \geq 4$ in W , there exists a shortcut s_{uvw} such that the edge multiset having representative vector $\chi^W + s_{uvw}$ is a walk with $t-1$ edges belonging to \mathcal{W}_c^- and containing the edge e . Necessarily, the edge (v, w) belongs to $\Delta_c(u)$.*

The sets $\Delta_c(u)$ may contain several edges; therefore, the application of Lemma 2.15 may require a lengthy procedure. The following lemma requires a weaker condition that is also easier to verify.

is almost Hamiltonian in w . We can assume that $k \geq 5$, since for $k = 3$ the edges (u_1^{i-2}, u_2^i) and (u^{i+1}, u_2^i) coincide. Consider the walk W_G^i obtained from W_F^i by replacing two edges of the cycle C^Y with two of its short chords, i.e., $W_G^i = W_F^i - (u_1^{i-1}, u_1^i) - (u_1^{i-2}, u_1^{i-3}) + (u_1^i, u_1^{i-2}) + (u_1^{i-1}, u_1^{i-3})$ (see Figure 12, where we used dotted and thick lines with the same convention as in Figure 11). From W_G^i we construct, as before, two Hamiltonian cycles to show that (u_1^{i-2}, u_2^i) and (u^{i+1}, u_2^i) are c -adjacent in w . It follows that the set J_w is c -connected in w .




Figure 12: The tight walk W_G^i built from W_F^i

For $w = u_{n_i}^i$, $i = 1, 2, \dots, k$, we set $J_w = \{(u_{n_{i-1}}^{i-1}, u_{n_{i-1}}^i), (u_{n_{i+1}}^{i+1}, u_{n_{i+1}}^i)\} \subseteq \Delta_c(w)$. The proof that in this cases the two edges of J_w are c -adjacent in w goes like for the cases $w = u_1^i$, $i = 1, 2, \dots, k$; for each i only one walk is used, actually the walk W_G^i taken “upside down”.



Figure 13: Example of reduction by shortcuts

It is easy to verify that each of the walks of the of the basis \mathcal{B}_c constructed in the proof of Theorem 2.12 can be reduced to an Hamiltonian cycle using the shortcuts defined by the sets J_w , $w \in V$. An example of such a reduction is shown in Figure 13 for the walk $W_3((u_1^1, u_1^4))$ of Figure 7(c). Again, for the dotted and the thick lines we follow the convention used for Figure 11. Observe how the sets J_w have three edges for w belonging to the cycle C^Y , while have only two edges for the nodes of the cycle C^Z . This is due to the fact that, for all the walks of \mathcal{B}_c , if a node w of C^Z has degree four, then the walk has a path edge and an edge of C^Z that are incident to w , which is not always the case if w belongs to C^Y .

Finally, by Theorem 2.12, all the walks of \mathcal{B}_c intersect the cycles C^Y and C^Z . It is easy to check that also the Hamiltonian cycles of \mathcal{C}_c obtained by shortcuts from those walks share this property. This proves that the properties (a) and (b) of the statement hold for \mathcal{C}_c . To prove that \mathcal{C}_c has also the properties (c) and (d), consider the walk $W_C^i(1)$ of Figure 5 for $i = 1, 2, \dots, k$. This walk contains the two cycle edges (u_1^i, u_1^{i-1}) and (u_1^i, u_1^{i+1}) of C^Y whose endnodes have both degree 2. Therefore, the Hamiltonian cycle of \mathcal{C}_c obtained by shortcuts from this walk also contains these two edges. Analogously, the Hamiltonian cycle of \mathcal{C}_c obtained from the walk $W_D^i(n_i - 1)$ contains both the cycle edges $(u_{n_i}^i, u_{n_{i-1}}^{i-1})$ and $(u_{n_i}^i, u_{n_{i+1}}^{i+1})$ of C^Z . \square

2.3. The 1-node lifting

We have seen in Section 2 that a simple bicycle inequality is the restriction of a simple wheelbarrow inequality obtained by removing the coefficients of the edges incident with the node Z . On the other hand, one can view a simple wheelbarrow as the extension of a simple bicycle inequality obtained by adding a node to its corresponding configuration. The operation of adding one node (and the edges incident with it) to the configuration of a generic facet defining inequality is called *1-node lifting* in [18], where the conditions on the coefficients of the new edges for the new inequality to preserve the face defining property are investigated. In particular, the theorem that follows states sufficient conditions for this to happen. When one of the edges incident with the new node has a zero coefficient, we have a special case of 1-node lifting, called *zero-lifting*, that is exploited in Section 3.

Theorem 2.18 (Theorem 4.4 [18]) *Let $cx \geq c_0$ be a TT inequality that is facet-defining for STSP(n); an inequality $c^*x^* \geq c_0$ which is obtained by 1-node lifting of $cx \geq c_0$ is facet-defining for STSP($n + 1$) if it is tight triangular and there exist an edge set $F \subseteq \Delta_{c^*}(u_{n+1})$ and a Hamiltonian cycle basis \mathcal{C}_c of $cx \geq c_0$ such that:*

- (i) $F \cap H \neq \emptyset$ for all $H \in \mathcal{C}_c$;
- (ii) for all $e \in F$ there exist $e' \neq e$, $e' \in \Delta_{c^*}(u_{n+1})$ and $H \in \mathcal{H}_c^-$ such that e and e' belong to H ;
- (iii) every connected component of the graph (V, F) contains at least one odd cycle;
- (iv) F is c^* -connected in u_{n+1} .

Remark 2.19. *Theorem 2.18 is proven in [18] by showing that a set of walks, obtained by first removing one or two edges of the set F from some walk of the canonical basis \mathcal{C}_c of $cx \geq c_0$, and then connecting the endnodes of the removed edges to the node u_{n+1} , contains a canonical basis \mathcal{C}_{c^*} of the inequality $c^*x^* \geq c_0$. Therefore, each walk of \mathcal{C}_{c^*} shares all the edges of some walk of \mathcal{C}_c , except some of those that intersect the set F .*

Using Theorem 2.18 and Remark 2.19 we can now show that the simple wheelbarrow inequalities define facets of STSP(n) and we can derive some properties of one of their canonical bases that will be used in the following to prove a similar theorem for the simple path inequalities.

Theorem 2.20. *A simple wheelbarrow inequality $cx \geq c_0$ associated with the simple k -wheelbarrow configuration $W(n_1, n_2, \dots, n_k)$ is facet defining for STSP(n) and has a Hamiltonian cycle basis \mathcal{C}_c with the following properties:*

- (a) Every Hamiltonian cycle of \mathcal{C}_c intersects the edge set of the cycle C_Z .
- (b) For each edge $e \in C_Z$ there exists $e' \in C_Z$ and a Hamiltonian cycle $H \in \mathcal{C}_c$ containing both e and e' .

Proof. A simple wheelbarrow inequality with the coefficients given in (11) is in TT form, since it is facet defining for GTSP(n) (see Proposition 2.2 [18]). As it has been already observed, it can be obtained by 1-node lifting of a bicycle inequality $\hat{c}\hat{x} \geq c_0$. To prove that it is facet defining for STSP(n) we use Theorem 2.18. Let the set F , required by that theorem, be defined as follows:

$$F = C_Y \cup \left\{ (u_1^i, u_j^{i+1}) \mid i \in \{1, 2, \dots, k\}, j \in \{2, \dots, n_{i+1}\} \right\}.$$

The set F belongs to $\Delta_c(Y)$ and satisfies the conditions (i), (ii), and (iii) due to the properties (a) and (c) of Theorem 2.17 and to the fact that the cycle C_Y has odd length. We only need

prove that F is c -connected in Y . For $i = 1, 2, \dots, k$, consider the walk $W_C^i(1)$ of Figure 5. Such a walk is tight for the bicycle inequality $\hat{c}\hat{x} \geq c_0$. This walk contains the two cycle edges $e = (u_1^i, u_1^{i-1})$ and $e' = (u_1^i, u_1^{i-2})$ of C_Y whose endnodes have both degree 2. Therefore, the Hamiltonian cycle $H \in \mathcal{H}_{\hat{c}}^-$, obtained by shortcuts from this walk, also contains these two edges. The walk obtained by removing e and e' from H and adding the four edges that connect the endnodes of the removed edges to Y is almost Hamiltonian in Y , belongs to \mathcal{W}_c^- , and implies that the two edges e and e' are c -adjacent in Y . Repeating this argument for the walk $W_D^{i-2}(1)$, we see that (u_1^{i-2}, u_1^{i-1}) and (u_1^i, u_1^{i-2}) are c -adjacent in Y as well. Consequently, the edge set C_Y is c -connected in Y . To show that any edge $e \in F \setminus C_Y$ is c -adjacent in Y to some edge of C_Y , it is sufficient to start with a Hamiltonian cycle $H \in \mathcal{H}_{\hat{c}}^-$ that contains e and an edge $e' \in C_Y$. Such a Hamiltonian cycle exists by Property (a) of Theorem 2.17. By removing e and e' from H and adding the edges that connect their endnodes to Y we produce a walk almost Hamiltonian in Y , implying that e and e' are c -adjacent in Y . Since the cycle C_Z has empty intersection with the edge set F , the property (a) and (b) of the Hamiltonian cycle basis \mathcal{C}_c follow from Remark 2.19 and the properties (b) and (d) of Theorem 2.17. \square

Theorem 2.21. *A simple path inequality $cx \geq c_0$ associated with the simple k -path configuration $P(n_1, n_2, \dots, n_k)$ is facet defining for STSP(n).*

Proof. A simple path inequality with the coefficients given in (11) is in TT form, since it is facet defining for GTSP(n) (see Proposition 2.2 [18]). In addition it can be obtained by 1-node lifting of a simple wheelbarrow inequality $\hat{c}\hat{x} \geq c_0$. To prove that it is facet defining for STSP(n) we use again Theorem 2.18. Let the set F required by that theorem be defined as follows:

$$F = C_Z \cup \left\{ (u_{n_i}^i, u_j^{i+1}) \mid i \in \{1, 2, \dots, k\}, j \in \{1, \dots, n_{i+1} - 1\} \right\} \cup \{(Y, u_{n_1}^1)\}.$$

The set F belongs to $\Delta_c(Z)$ and satisfies the conditions (i), (ii), and (iii) due to the properties (a) and (b) of Theorem 2.20 and to the fact that the cycle C_Z has odd length. We only need prove that F is c -connected in Z . For $i = 1, 2, \dots, k$, consider the walk obtained from $W_D^i(n_i - 1)$ of Figure 5 by deleting an edge belonging to C_Y and connecting its endnodes to Y . Such a walk is tight for the simple wheelbarrow inequality $\hat{c}\hat{x} \geq c_0$. This walk contains the two cycle edges $e = (u_{n_i}^i, u_{n_{i-1}}^{i-1})$ and $e' = (u_{n_i}^i, u_{n_{i+1}}^{i+1})$ of C_Z whose endnodes have both degree 2. Therefore, the Hamiltonian cycle $H \in \mathcal{H}_{\hat{c}}^-$, obtained by shortcuts from this walk, also contains these two edges. The walk obtained by removing e and e' from H and adding the four edges that connect the endnodes of the removed edges to Z is almost Hamiltonian in Z , belongs to \mathcal{W}_c^- , and implies that the two edges e and e' are c -adjacent in Z . Consequently, C_Z is c -connected in Z . To show that any edge $e \in F \setminus C_Z$ and some edge of C_Y are c -adjacent in Z , it is sufficient to start with a Hamiltonian cycle $H \in \mathcal{H}_{\hat{c}}^-$ that contains e and an edge $e' \in C_Z$. Such a Hamiltonian cycle exists by Property (a) of Theorem 2.20. By removing e and e' from H and adding the edges that connect their endnodes to Z , we produce a walk almost Hamiltonian in Z that implies that e and e' are c -adjacent in Z . \square

The theorems 2.17, 2.20, and 2.21 are summarized in the following.

Theorem 2.22. *The simple PWB inequalities are facet defining for STSP(n).*

3. The PWB inequalities

As we have seen in Section 1, the configuration of a PWB inequality $c^*x \geq c_0$ is obtained from the configuration of a simple one $cx \geq c_0$ (same right hand side), by replacing each nodes i with

a node set V^i of arbitrary size. As the edges in $\gamma(V^i)$ have a zero coefficient, the operation of replacing a node i by a set V^i in a configuration is called *zero-lifting* of i .

The following theorem, proved in [18], gives a sufficient condition to derive the facet-defining property of an inequality from the facet-defining property of its simple archetype.

Theorem 3.1 (Theorem 4.9 [18]) *A TT inequality $cx \geq c_0$ is facet-defining for STSP(n) if its associated simple inequality $\bar{c}\bar{x} \geq c_0$ is nontrivial and facet-defining for STSP(p) and for every $v \in V_p$ the set $\delta(v)$ in K_p is \bar{c} -connected.*

We first apply this theorem to the bicycle inequalities.

Theorem 3.2. *The bicycle inequalities are facet defining for STSP(n).*

Proof. Let $c^*x^* \geq c_0$ be a bicycle inequality and $cx \geq c_0$ be its associated simple bicycle inequality. By converting the walks of the figures 5, 7, and 8 to Hamiltonian cycles using shortcuts, it is easy to verify the following:

(a) For $i = 1, 2, \dots, k$ and for each edge e incident with u_1^i with $e \neq (u_1^i, u_2^i)$, there exists a Hamiltonian cycle of \mathcal{H}_c^- containing both e and (u_1^i, u_2^i) .

(b) For $i = 1, 2, \dots, k$ and for each edge e incident with $u_{n_i}^i$ with $e \neq (u_{n_i-1}^i, u_{n_i}^i)$, there exists a Hamiltonian cycle of \mathcal{H}_c^- containing both e and $(u_{n_i-1}^i, u_{n_i}^i)$.

(c) For $i = 1, 2, \dots, k$ with $n_i \geq 3$ and for each edge e incident with u_j^i with $1 < j < n_i$, $e \neq (u_j^i, u_{j-1}^i)$, and $e \neq (u_j^i, u_{j+1}^i)$, there exists a Hamiltonian cycle of \mathcal{H}_c^- containing either the pair of edges e and (u_j^i, u_{j-1}^i) or the pair e and (u_j^i, u_{j+1}^i) .

(d) For $i = 1, 2, \dots, k$ with $n_i \geq 3$, there exists a Hamiltonian cycle of \mathcal{H}_c^- containing both (u_j^i, u_{j-1}^i) and (u_j^i, u_{j+1}^i) .

From (a), (b), (c), and (d) it follows that $\delta(u_j^i)$ is c -connected for $i = 1, 2, \dots, k$ and for $j = 1, \dots, n_i$. Therefore, by Theorem 3.1, the theorem follows. \square

Now we make use of the following lemma to extend the facet-defining property to all the PWB inequalities.

Lemma 3.3 (Lemma 4.8 [18]) *Let $cx \geq c_0$ be a TT inequality facet-defining for STSP(n), $c^*x^* \geq c_0$ be an inequality obtained by 1-node lifting of $cx \geq c_0$ and F be a subset of $\Delta_{c^*}(u_{n+1})$ that satisfies the conditions of Theorem 2.18. Then the following hold:*

(a) *The edge set $\delta(u_{n+1}) \subseteq E_{n+1}$ is c^* -connected if the graph (V, F) is connected.*

(b) *For every $v \in V$ the edge set $\delta(v) \subseteq E_{n+1}$ is c^* -connected if the edge set $\delta(v) \subseteq E_n$ is c -connected.*

Theorem 3.4. *The PWB inequalities are facet defining for STSP(n).*

Proof. It follows from Theorem 3.2, from Lemma 3.3, and from the connectivity of the graph induced by the edge set F defined in the theorems 2.20 and 2.21. \square

Many of the inequalities known to define facets of STSP(n) are special cases of PWB inequalities, as it can be easily verified by putting them in TT form. In particular,

(i) the *2-matching* inequalities, introduced by Edmonds, are 2-regular PWB inequalities obtained from a simple inequality possibly by zero-lifting of the nodes Y and Z ;

(ii) the *Chvátal comb* inequalities, introduced by Chvátal, are 2-regular PWB inequalities obtained from a simple inequality possibly by zero-lifting of Y , Z , and of the nodes of the cycle C_Y ;

(iii) the *comb* inequalities, introduced by Grötschel and Padberg, are 2-regular PWB inequalities.

4. The extended PWB inequalities

Some simple inequalities can be obtained by extending other simple inequalities by simultaneously adding two nodes to their configuration. An example of such an extension is called *edge cloning* in [18] and it is defined as follows:

Definition 4.1. Let $cx \geq c_0$ be a *TT* inequality defined on \mathbb{R}^{E_n} and e be an edge in E_n . We say that the inequality $c^*x^* \geq c_0^*$ defined on $\mathbb{R}^{E_{n+2t}}$, with $t \geq 1$, is obtained from $cx \geq c_0$ by cloning t times edge $e = (u_{n-1}, u_n)$ if (see Figure 14, in which $t = 1$):

$$\begin{aligned} c_0^* &= c_0 + 2tc_e \\ c^*(u_i, u_{n+j}) &= c(u_i, u_{n-1}) \quad \text{for } 1 \leq i \leq n-2, 1 \leq j \leq 2t-1 \text{ and } j \text{ odd,} \\ c^*(u_i, u_{n+j}) &= c(u_i, u_n) \quad \text{for } 1 \leq i \leq n-2, 2 \leq j \leq 2t \text{ and } j \text{ even,} \\ c^*(u_{n+i}, u_{n+j}) &= 2c_e \quad \text{for } -1 \leq i < j \leq 2t \text{ and } j-i \text{ even,} \\ c^*(u_{n+i}, u_{n+j}) &= c_e \quad \text{for } -1 \leq i < j \leq 2t \text{ and } j-i \text{ odd.} \end{aligned}$$

The nodes u_{n+j} for $1 \leq j < 2t$ and j odd and the nodes u_{n+j} for $1 < j \leq 2t$ and j even are called the clones of the nodes u_{n-1} and u_n , respectively. The edge $e = (u_{n-1}, u_n)$ is said to be cloned.




Figure 14: Cloning edge (u_{n-1}, u_n) one time

Let $cx \geq c_0$ be a *TT* inequality defining a facet of $STSP(n)$; an edge $e = (u, v)$ is called *c-clonable* if the c -length of every walk W of K_n is at least $c_0 + (d_e(W) - 2)c_e$, where $d_e(W)$ is the minimum of the degrees of u and v in W ; we say that a node $v \in V$ is α -critical for the inequality if the c -length of a minimum c -length walk of $K_n - \{v\}$ is $c_0 - \alpha$.

Theorem 4.2 (Theorems 4.12 and 4.13 [18]) Let $cx \geq c_0$ be a nontrivial *TT* inequality facet-defining for $STSP(n)$ and $e = (u_{n-1}, u_n)$ be a c -clonable edge such that u_{n-1} and u_n are $2c_e$ -critical for $cx \geq c_0$. Then the following hold:

- (a) the inequality $c^*x^* \geq c_0^*$ obtained by cloning e (t times) is facet-defining for $STSP(n+2t)$;
- (b) the edge subsets $\delta(u_{n-1}), \dots, \delta(u_{n+2t})$ of E_{n+2t} are c^* -connected;
- (c) for $v \in V - \{u_{n-1}, u_n\}$, if $\delta(v)$ in K_n is c -connected, then $\delta(v)$ in K_{n+2t} is c^* -connected;
- (d) if $f = (z_1, z_2) \neq e$ is an edge in E_n such that z_1 and z_2 are $2c_f$ -critical for $cx \geq c_0$, then z_1 and z_2 are $2c_f^*$ -critical for $c^*x \geq c_0^*$.

Definition 4.3. A simple extended PWB inequality is the inequality obtained by cloning t_i times the edge (u_1^i, u_2^i) for each $i \in I$ where I is any subset of $\{1, \dots, k\}$ such that $n_i = 2$ for all $i \in I$. An inequality is an extended PWB if its associated simple inequality is a simple extended PWB.

Theorem 4.4. *The extended PWB inequalities are facet defining for STSP(n).*

Proof. Let $cx \geq c_0$ be a simple PWB inequality of STSP(n) with $n_i = 2$ for some i . We first prove the following two claims.

Claim 4.5. *Let $e = (u_1^i, u_2^i)$, with $n_i = 2$, be an edge of an extended PWB configuration that is not cloned and let $cx \geq c_0$ be the inequality associated with the configuration. Then the validity of $cx \geq c_0$ implies that e is c -clonable.*

Proof. We have to show that if e is not c -clonable, i.e., if there exists a walk W of K_n such that $c(W) < c_0 - (d_e(W) - 2)c_e$, then the inequality is not valid, i.e., there exists a walk W^* with $c(W^*) < c_0$. If $d_e(W) = 2$, then obviously $W^* = W$. If $d_e(W) \geq 4$ we show how to construct a walk W' from W where the minimum degree of u_1^i and u_2^i is $d_e(W) - 2$ and $c(W') \leq c(W) - 2c_e$. Thus, we can construct the walk W^* by recursively applying this process. We consider three distinct cases.

Case (a). W contains at least three copies of e . The walk W' is obtained from W by removing two copies of e . The multigraph $K_n[W']$ is connected and $c(W') = c(W) - 2c_e$.

Case (b). W contains two copies of e . Let $u_r^q \notin \{u_1^i, u_2^i\}$ and $u_t^s \notin \{u_1^i, u_2^i\}$ be neighbors in W of u_1^i and u_2^i , respectively, such that the multigraph $K_n[W']$ is connected, where $W' = W - e - (u_1^i, u_r^q) - (u_2^i, u_t^s) + (u_r^q, u_t^s)$. Since $n_i = 2$, it follows that either $u_r^q \in \{Y, Z\}$ or $q \neq i$ and either $u_t^s \in \{Y, Z\}$ or $s \neq i$. Thus, it is easy to verify from (11) that $c(W') \leq c(W) - 2c_e$ and that equality holds only if $u_r^q = Y$ and $u_t^s = Z$.

Case (c). W contains only one copy of e . Among the edges of W different from e that are incident with u_1^i , there are always two, say (u_1^i, w) and (u_1^i, z) , or, possibly, two copies of the same edge (in which case $w = z$), such that $K_n[W'']$ is connected, where $W'' = W - (u_1^i, w) - (u_1^i, z) + (w, z)$. Specularly, among the edges of W different from e that are incident with u_2^i , there are always two, say (u_2^i, w') and (u_2^i, z') , or, possibly, two copies of the same edge (in which case $w' = z'$), such that $K_n[W']$ is connected, where $W' = W'' - (u_2^i, w') - (u_2^i, z') + (w', z')$. It is easy to verify from (11) that $c(W') \leq c(W) - 2c_e$. \square

Observe that if $n_i > 2$, then none of the edges of the path π^i is c -clonable, because there are tight walks where both the endnodes of a path edge have degree four (see, e.g., Figure 5). This explains the exclusion of these edges from the cloning process in Definition 4.3.

Claim 4.6. *Both the endnodes of the edge $e = (u_1^i, u_2^i)$ are $2c_e$ -critical.*

Proof. The walk W_A^i of Definition 2.10 contains two copies of e . The Hamiltonian cycle H_1 of $K_n - \{u_1^i\}$, obtained from W_A^i by removing the two copies of e , has length $c_0 - 2c_e$. It follows that u_1^i is $2c_e$ -critical. Analogously, using the walk W_B^i , one proves that also u_2^i is $2c_e$ -critical. \square

Let ℓ be the number of edges that have been cloned to obtain the inequality $cx \geq c_0$, i.e., $\ell = |\{i \mid t_i > 0\}|$. If $\ell = 0$, the inequality is valid since it is a simple PWB. Therefore, by Claim 4.5, $e = (u_1^i, u_2^i)$ with $n_i = 2$ is c -clonable and, by Claim 4.6 both its endnodes are $2c_e$ -critical. Thus, applying Theorem 4.2, the inequality $c^*x^* \geq c_0^*$, obtained from $cx \geq c_0$ by cloning e for t_i times, is facet defining for STSP($n + 2t_i$) and, for any node v , the edge set $\delta(v)$ in K_{n+2t_i} is c^* -connected. In addition, if the endnodes of an edge f are $2c_f$ -critical, then they are also $2c_f^*$ -critical. Finally, by induction on ℓ one shows that any simple extended PWB inequality $\hat{c}x \geq \hat{c}_0$ is facet defining and that all its nodes are \hat{c} -connected. Thus, any extended PWB inequality is facet defining. \square

In [24] Padberg and Hong define the *chain inequality* as follows:

Let $S_i \subseteq V$ for $i = 0, 1, \dots, q$ be any proper subsets of V satisfying $S_i \cap S_0 = \emptyset$ for $i = 1, \dots, p$, $|S_i \cap S_0| \geq 1$ and $|S_i \setminus S_0| \geq 1$ for $i = p+1, \dots, q$, and $S_i \cap S_j = \emptyset$ for $1 \leq i < j \leq q$. Let $R \subseteq S_0$ be a subset of S_0 satisfying $|R| = p$ and $R \cap S_i = \emptyset$ for $i = 1, \dots, q$. Then the *chain inequality* is

$$\sum_{i=0}^q x(\gamma(S_i)) + \sum_{i=1}^p x((R : S_i)) \leq |S_0| + |R| + \sum_{i=1}^q (|S_i| - 1) - \left\lfloor \frac{q-p+1}{2} \right\rfloor. \quad (22)$$

In [24] it is shown that the chain inequality is a valid inequality for STSP(n) with $n \geq 8$, when $2 \leq p < q$ (for $p \leq 1$ the chain inequality coincides with a comb inequality). If $q - p$ is odd, then the inequality is dominated by a nonnegative linear combination of subtour elimination and degree constraints (see, e.g., [15] where a generalization of these inequalities is described).

It is not difficult to verify, by putting (22) in TT form, that the chain inequality is an extended 2-regular PWB inequality obtained by cloning only one path edge of a simple 2-regular PWB configuration, say (u_1^1, u_2^1) , and then zero-lifting all nodes except u_1^1 and its clones (these are the members of the set R in the definition of the chain inequalities).

Extended PWB inequalities generalize chain inequalities not only because the cloning process can involve all path edges of a 2-regular PWB inequality and because all nodes can be zero-lifted, but also because these inequalities can be derived from any non-regular PWB inequality having at least one path of length 2. This generalization of the chain inequalities differs from the one described in [15].

5. Composition of PWB inequalities

In [17] an operation called *2-sum* is described, which yields facet defining inequalities for GTSP(n) and involves two facet defining inequalities defined on smaller configurations. In [18] sufficient conditions are given under which an inequality, obtained as the 2-sum of two other inequalities, defines a facet of STSP(n). We apply the 2-sum operation to PWB inequalities. For the sake of completeness, we give the formal definition of the 2-sum operation.

Two weighted graphs $G^1 = (V^1, E^1, c^1)$ and $G^2 = (V^2, E^2, c^2)$ are *isomorphic* if there exists a one-to-one correspondence ρ between their node sets that preserves the weight function, i.e., for every edge $(u, v) \in E^1$, the edge $(\rho(u), \rho(v))$ belongs to E^2 and $c^1(u, v) = c^2(\rho(u), \rho(v))$.

Definition 5.1. Let $c^1 x^1 \geq c_0^1$ and $c^2 x^2 \geq c_0^2$ be two TT inequalities facet defining for STSP(n_1) and STSP(n_2), respectively, and let $e_1 = (u_1, v_1) \in E_{n_1}$ and $e_2 = (u_2, v_2) \in E_{n_2}$ be two edges such that $c^1(u_1, v_1) = c^2(u_2, v_2) = \theta > 0$. Denote by V_{n_1} and V_{n_2} the node sets of the two graphs K_{n_1} and K_{n_2} , respectively, by V^1 the set $V_{n_1} - \{u_1, v_1\}$ and by V^2 the set $V_{n_2} - \{u_2, v_2\}$. Then the 2-sum of the two inequalities, obtained by identifying u_1 with u_2 and v_1 with v_2 , is the inequality $cx \geq c_0^1 + c_0^2 - 2\theta$ defined on \mathbb{R}^{E_n} , with $n = n_1 + n_2 - 2$, whose support graph $G_c = (V, E_n, c)$ is defined as follows:

- (i) $V = V^1 + V^2 + \{u, v\}$;
- (ii) the subgraph of G_c induced by $V^1 + \{u, v\}$ is isomorphic to G_{c^1} and u and v correspond to u_1 and v_1 , respectively, in the isomorphism;
- (iii) the subgraph of G_c induced by $V^2 + \{u, v\}$ is isomorphic to G_{c^2} and u and v correspond to u_2 and v_2 , respectively, in the isomorphism;
- (iv) the coefficients of the edges with one endnode in V_1 and the other in V_2 , that we call the crossing edges of the 2-sum, are computed by a sequential lifting procedure.

The inequalities $c^1 x^1 \geq c_0^1$ and $c^2 x^2 \geq c_0^2$ are called the *component inequalities* of the 2-sum.

For the sake of simplicity, from now on every time that the correspondence between nodes, edges, and walks of each of the graph G_{c_1} and G_{c_2} and their corresponding isomorphic subgraphs of G_c is evident, we omit to mention it explicitly.

In [17] the 2-sum operation is used in a recursive way to generate huge families of inequalities defining facets of GTSP(n). The most interesting of such families is perhaps the one of the *regular parity path tree* (or *regular parity PWB-tree*) inequalities. These inequalities are peculiar because the lifting procedure does not have to be carried out explicitly, since the coefficient of each crossing edge is given by the c -length of the shortest path between its endnodes in the c -weighted graph produced by each 2-sum. This implies that the coefficients of the crossing edges do not depend on the order of the sequential lifting procedure.

Definition 5.2. *A regular parity PWB-tree inequality is one of the following:*

(i) *A regular PWB inequality \mathcal{P}_0 . In this case the PWB-tree inequality has length 0. The path edges as well as the even and the odd nodes of \mathcal{P}_0 coincide with the corresponding ones of the PWB inequality.*

(ii) *The 2-sum $\mathcal{P}_{\ell+1}$ of a regular parity PWB-tree inequality \mathcal{P}_ℓ , having length ℓ , and of a regular PWB inequality obtained by identifying the endnodes of a path edge of one inequality with the endnodes of a path edge of the other, with the constraints that the identified nodes have the same parity. In this case the PWB-tree inequality has length $\ell+1$. The nodes that result from the two identifications inherit the parity of the nodes from which they derive; the edge between them is a path edge of $\mathcal{P}_{\ell+1}$. The path edges as well as the even and the odd nodes of \mathcal{P}_ℓ and of the PWB inequality are path edges, even and odd nodes of $\mathcal{P}_{\ell+1}$.*

As it was mentioned before, for the regular parity PWB-tree inequalities the following theorem holds:

Theorem 5.3 (Theorem 5.3 [17]) *Let $c^1x^1 \geq c_0^1$ and $c^2x^2 \geq c_0^2$ be two regular parity PWB-tree inequalities, $cx \geq c_0$ be their 2-sum, and (u, v) be the edge that results from the identification. Then, using the notation of Definition 5.1, for any $x \in V^1$ and $y \in V^2$ the following holds:*

$$c(x, y) = \min \{c^1(x, u) + c^2(u, y), c^1(x, v) + c^2(v, y)\}. \quad (23)$$

The main result of this section is a proof that regular parity PWB-tree inequalities are facet defining for STSP(n). To do so, we use the following theorem that gives conditions for a 2-sum inequality to be facet defining for STSP(n).

We call a 2-sum inequality *h-liftable* if the coefficients of its crossing edges do not change if Hamiltonian cycles are used in the lifting procedure, instead of walks.

Theorem 5.4 (Theorem 3.5 [18]) *Under the assumptions of Definition 5.1, let $c^1x^1 \geq c_0^1$ and $c^2x^2 \geq c_0^2$ be nontrivial inequalities defining facets of STSP(n_1) and STSP(n_2), respectively. The 2-sum inequality $cx \geq c_0$ is facet-defining for STSP(n) if it is h-liftable and:*

(a) v_1 is 2θ -critical for $c^1x^1 \geq c_0^1$,

(b) $\delta(u_2)$ is c^2 -connected,

and either (Case (A))

(c') u_2 is 2θ -critical for $c^2x^2 \geq c_0^2$,

(d') $\delta(v_1)$ is c^1 -connected,

or (Case (B))

(c'') v_2 is 2θ -critical for $c^2x^2 \geq c_0^2$,

(d'') $\delta(u_1)$ is c^1 -connected,

(e'') there exists a Hamiltonian cycle $H_1 \in \mathcal{H}_{c_1}^-$ containing the edge (u_1, v_1) and any edge $e_1 \in \Delta_{c_1}(v_1)$,

(f'') there exists a Hamiltonian cycle $H_2 \in \mathcal{H}_{c_2}^-$ containing the edge (u_2, v_2) and any edge $e_2 \in \Delta_{c_2}(v_2)$.

Before stating the main theorem, we prove a lemma concerning the h -liftability of simple regular parity PWB-tree inequalities and recall three lemmata from [18] that will be used in the main proof as well.




Figure 15: Tight walks containing edge (u_j^i, u_q^r) and twice edge (u_j^i, u_{j+1}^i)

Lemma 5.5. *The 2-sum of a simple regular parity PWB-tree and of a simple PWB inequality is h -liftable.*

Proof. We consider the 2-sum of a simple regular parity PWB-tree inequality \mathcal{P}_ℓ with a simple regular PWB inequality $cx \geq c_0$, obtained by identifying the endnodes of a path edge of \mathcal{P}_ℓ with the endnodes of the edge (u_j^i, u_{j+1}^i) of the PWB inequality. Without loss of generality, we can assume that $j < n_i$, and thus that u_{j+1}^i is always even. Let $e = (a, u_q^r)$ be a crossing edge of the 2-sum inequality. We consider two cases, depending on whether node u_j^i (as well as the corresponding node of \mathcal{P}_ℓ) is even or odd. We use the walks of the figures 15, 16, and 17. These walks belong to \mathcal{W}_c^- when $cx \geq c_0$ is a path inequality and can be easily adjusted to the cases when the inequality is either a wheelbarrow or a bicycle.

Case A. Node u_j^i is even. Without loss of generality, we assume that $q \leq j$ (else, just exchange the roles of j and $j+1$ and of Z and Y). Then it is easy to see, by (23) and (13), that $c_e = c^1(a, u_j^i) + c^2(u_j^i, u_q^r)$. The inequality \mathcal{P}_ℓ is facet defining for GTSP(n) (see [17]). Therefore, there exists a tight walk containing (a, u_j^i) that, by Lemma 2.15, can be turned into a Hamiltonian cycle H containing (a, u_j^i) . If $r = i$, then let W_1 be the walk of Figure 15(a); otherwise, let W_1 be the walk of Figure 15(b). This walk contains edge (u_j^i, u_{j+1}^i) twice. The Hamiltonian cycle $H + W_1 - 2(u_j^i, u_{j+1}^i) - (a, u_j^i) - (u_j^i, u_q^r) + (a, u_q^r)$ is tight for the 2-sum inequality and contains the crossing edge e .

Case B. Node u_j^i is odd, i.e., $u_j^i = Y$ and $j = 0$. In [17] (Lemma 5.6) it is shown that there exists a tight walk W for the inequality \mathcal{P}_ℓ that contains both edges (a, Y) and (Y, u_1^i) . We




Figure 16: Tight walks containing the edges (u_1^i, Y) and (Y, u_q^r)




Figure 17: A tight walk containing the edges (u_1^i, Y) and (Y, u_q^r) and one containing twice edge (u_1^i, Y) and edge (u_1^i, u_q^r)

first assume that either $r \neq i$ or $q = n_r + 1$, in the latter case we have $u_q^r = Z$. Then it is easy to see that, by (23) and (11), $c_e = c^1(a, Y) + c^2(Y, u_q^r)$. If $q = n_r + 1$, then let W_2 be the walk of Figure 16(a); otherwise let it be the walk of Figure 16(b) (the edges represented by broken lines have to replace the marked edge if node Y has degree 2 in W). The walk $W + W_2 - 2(u_1^i, Y) - (a, Y) - (Y, u_q^r) + (a, u_q^r)$ is tight for the 2-sum inequality, contains the crossing edge e , and, by Lemma 2.15, can be transformed into a tight Hamiltonian cycle by shortcuts involving only edges of E^1 since only one edge on E^2 is incident with u_1^i and Y , respectively. Now let us assume that $r = i$ and $q > 0$. If $c_e = c^1(a, Z) + c^2(Z, u_q^r)$, then let W_3 be the walk of Figure 17(a). The walk $W' = W + W_3 - 2(u_1^i, Y) - (a, Y) - (Y, u_q^r) + (a, u_q^r)$ is tight for the 2-sum inequality and contains the crossing edge e . Finally, If $c_e = c^1(a, u_1^i) + c^2(u_1^i, u_q^r)$, let H be the Hamiltonian cycle defined for Case (A) and W_4 the walk of Figure 17(b). Then also the walk $W'' = H + W_4 - 2(u_1^i, Y) - (a, u_1^i) - (u_1^i, u_q^r) + (a, u_q^r)$ is tight for the 2-sum and contains e . Both walks W' and W'' can be turned into tight walks where every node but Y has degree 2, by applying shortcuts involving only edges of E^1 , since only one edge on E^2 is

incident with u_1^i . To reduce each of these walks to a Hamiltonian cycle, a shortcut involving a crossing edge (w, u_t^s) , where $s \neq i$, is necessary. However, since the coefficients of the inequality do not depend on the lifting sequence, we can assume, without loss of generality, that the lifting coefficients for all the crossing edges (w, u_t^s) , with $s \neq i$, are computed first, and the lemma is proved. \square

The following three lemmata give conditions under which some properties of the components of a 2-sum are carried over to the resulting inequality. The notation used in their statements is that of Definition 5.1.

Lemma 5.6 (Lemma 3.5 [18]) *Under the assumptions of Definition 5.1, let $w \in V_{n_1}$ be α -critical for $c^1x^1 \geq c_0^1$, and $c_e^1 = \alpha/2$ for some edge $e \in \delta(w)$ in K_{n_1} . The corresponding node $w \in V$ is α -critical for $cx \geq c_0$ if $cx \geq c_0$ is supporting for GTSP(n) and any of the following conditions holds:*

- (a) $w \notin \{u_1, v_1\}$,
- (b) $w = u_1$ and u_2 is 2θ -critical for $c^2x^2 \geq c_0^2$,
- (c) $w = v_1$ and v_2 is 2θ -critical for $c^2x^2 \geq c_0^2$.

Lemma 5.7 (Lemma 3.6 [18]) *Under the assumptions of Definition 5.1, if there exists a Hamiltonian cycle $H \in \mathcal{H}_{c_1}^-$ containing two nonadjacent edges e and $f \in E_{n_1}$ and at least one of the two nodes u_2 and v_2 is 2θ -critical for $c^2x^2 \geq c_0^2$, then there exists a Hamiltonian cycle $H^* \in \mathcal{H}_c^-$ containing the edges in E_n corresponding to e and f , respectively.*

Lemma 5.8 (Lemma 3.7 [18]) *Under the assumptions of Definition 5.1, let $\delta(w)$ in K_{n_1} be c^1 -connected for every node $w \in V_{n_1}$ and let $\delta(w)$ in K_{n_2} be c^2 -connected for every node $w \in V_{n_2}$. Then $\delta(w)$ in K_n is c -connected for every node $w \in V$ if the following conditions hold:*

- (i) $cx \geq c_0$ is h -liftable;
- (ii) at least one of the two nodes u_1 and v_1 is 2θ -critical for $c^1x^1 \geq c_0^1$;
- (iii) at least one of the two nodes u_2 and v_2 is 2θ -critical for $c^2x^2 \geq c_0^2$.

We have now all the basic tools to prove the following

Theorem 5.9. *The regular parity PWB-tree inequalities are facet defining for STSP(n).*

Proof. Let \mathcal{P}_ℓ be a simple regular parity PWB-tree inequality of length ℓ . We proceed by induction on ℓ . The theorem is true for $\ell = 0$ since \mathcal{P}_0 is a PWB inequality. We consider now the case $\ell = 1$. Thus, by Definition 5.2, \mathcal{P}_1 is the 2-sum of an inequality \mathcal{P}_0 , which we denote by $c^1x^1 \geq c_0^1$, and of a simple PWB inequality $c^2x^2 \geq c_0^2$, obtained by identifying the endnodes of the path edge (u_1, v_1) of the first inequality to the endnodes of the path edge (u_2, v_2) of the second. Using the notation of Definition 5.1, we show now that the conditions of Theorem 5.4 are satisfied. By Lemma 5.5, \mathcal{P}_ℓ is h -liftable for all values of ℓ .

Claim 5.10. *An even endnode of a path edge e of a simple PWB inequality $cx \geq c_0$ is $2c_e$ -critical.*

Proof. Without loss of generality, we can assume that $e = (u_j^i, u_{j+1}^i)$, with $j \in \{0, \dots, n_i - 1\}$. We consider two cases.

The first case arises when $j \in \{1, \dots, n_i - 1\}$; in this case both the endnodes of e are even. Consider the walk W_A^i of Definition 2.10 if $j = 1$, the walk W_B^i if $j = n_i - 1$, and $W_C^i(j + 1)$ if

$j \in \{2, \dots, n_i - 2\}$. Such a walk contains two copies of edge e ; in addition, by removing these two copies, we obtain a walk of $V \setminus \{u_{j+1}^i\}$ of c -length $c_0 - 2c_e$. Therefore, u_{j+1}^i is $2c_e$ -connected. Similarly, one shows that also u_j^i is $2c_e$ -connected.

In the second case, arising when $j = 0$, we consider the walk $W_C^i(1)$, the only even node is u_1^i . We add the (odd) node $Y = u_0^i$, then we remove the two cycle edges (u_1^i, u_1^{i-1}) and (u_1^i, u_1^{i+1}) , and finally we add the edges (Y, u_1^{i-1}) , (Y, u_1^{i+1}) , and two copies of the edge $e = (Y, u_1^i)$. The resulting walk has c -length c_0 and contains two copies of e ; moreover, the removal of these two copies of e yields a walk of $V \setminus \{u_j^i\}$ of c -length $c_0 - 2c_e$. \square

Claim 5.11. *An even endnode of a path edge e of a simple regular parity PWB-tree inequality $cx \geq c_0$ is $2c_e$ -critical.*

Proof. It follows immediately by applying the recursion of Definition 5.2, Claim 5.11, and Lemma 5.6. \square

Since a path edge has at most one odd endnode, we can assume, without loss of generality, that the nodes u_1 and u_2 are both even. Thus, by claims 5.10 and 5.11, the conditions (a) and (c'') are satisfied for all values of ℓ . Depending on the parity of the nodes v^1 and v^2 we have two distinct cases.

Case A: v^1 and v^2 are even. In this case, by claims 5.10 and 5.11 Condition (c') is satisfied.

Case B: v^1 and v^2 are odd. First, we prove the following

Claim 5.12. *Let $cx \geq c_0$ be a simple PWB inequality and (u_1^i, Y) be one of its path edges incident with an odd node. Then there exists a Hamiltonian cycle $H \in \mathcal{H}_c^-$ containing (u_1^i, Y) and an edge of $\Delta_c(Y)$.*

Proof. Consider the walk of Figure 7(a), where j and q are set to 1 and n_r , respectively. With these settings of j and q the walk is actually a Hamiltonian cycle. Replace $(u_1^i, u_{n_r}^r)$ by the two edges (u_1^i, Y) and $(Y, u_{n_r}^r)$. For the case of a path inequality, insert node Z into the cycle in a similar manner, by removing any cycle edge of C_Z (by Theorem 2.12 at least one edge of C_Z belongs to the Hamiltonian cycle). The resulting Hamiltonian cycle belongs to \mathcal{H}_c^- , contains the edge (u_1^i, Y) , and, again by Theorem 2.12, intersects $C_Y \subseteq \Delta_c(Y)$. \square

Claim 5.13. *Let $cx \geq c_0$ be a simple regular parity PWB-tree inequality and (u, v) be one of its path edges incident with an odd node v . Then there exists a Hamiltonian cycle $H \in \mathcal{H}_c^-$ containing (u, v) and an edge of $\Delta_c(v)$.*

Proof. It follows immediately by applying the recursion of Definition 5.2, Claim 5.10, and Lemma 5.7. \square

End of proof of Theorem 5.9. Claims 5.12 and 5.13 imply that both conditions (e'') and (f'') are satisfied for all values of ℓ .

For $\ell = 1$ the two inequalities are of PWB type, hence, by theorems 3.2 and 3.4, the conditions (b), (d'), and (d'') are satisfied. Thus, the inequality is facet defining and each node of the graph is c -connected with respect to the coefficient vector c of the inequality. This implies that the same holds for $\ell > 1$, by applying the induction and the lemmata 5.5 and 5.8. \square

A clique-tree inequality (see [11]), as long as there is a node not contained in any handle or tooth, is a special case of a regular parity path three obtained as follows (see [17]):

- only 2-regular PWB inequalities with at least one odd node are used as components,

- each 2-sum involves an odd node,
- all odd nodes involved in the 2-sums correspond to a single odd node in the resulting inequality.

It is evident how the regular parity PWB-tree inequalities generalize the clique tree inequalities in many possible ways.

6. Conclusions

The operations described in this paper can be extended in several directions to prove that further classes of inequalities derived from PWB inequalities are facet defining for the STSP polytope. For example, the components of a 2-sum can be extended PWB inequalities, or edge cloning can be applied to regular parity PWB inequalities.

References

- [1] E. Balas and M. Fischetti, “On the monotonization of polyhedra,” *Mathematical Programming*, vol. 77, pp. 59–84, 1997.
- [2] T. Christof and G. Reinelt, “Combinatorial optimization and small polytopes,” *TOP*, vol. 4, pp. 1–53, 1996.
- [3] V. Chvátal, “Edmonds polytopes and weakly hamiltonian graphs,” *Mathematical Programming*, vol. 5, pp. 29–40, 1973.
- [4] J. Clochard and D. Naddef, “Using path inequalities in a branch and cut code for the symmetric traveling salesman problem,” in *Proceedings of the 3rd Mathematical Programming Conference on Integer Programming and Combinatorial Optimization* (G. Rinaldi and L. Wolsey, eds.), (Louvain-la-Neuve, Belgium), pp. 291–311, CORE, 1993.
- [5] G. Cornuéjols, J. Fonlupt, and D. Naddef, “The traveling salesman problem on a graph and some related integer polyhedra,” *Mathematical Programming*, vol. 33, pp. 1–27, 1985.
- [6] M. Goemans, “Worst-case comparison of valid inequalities for the TSP,” *Mathematical Programming*, vol. 69, pp. 335–349, 1995.
- [7] M. Grötschel, “On the monotone symmetric travelling salesman problem: hypohamiltonian/hypotractable graphs and facets,” *Mathematics of Operations Research*, vol. 5, pp. 285–292, 1980.
- [8] M. Grötschel and L. Lovász, “Combinatorial Optimization,” in *Handbook of Combinatorics* (R. Graham, M. Grötschel, and L. Lovász, eds.), vol. II, ch. 28, pp. 1541–1597, Amsterdam: North-Holland, 1995.
- [9] M. Grötschel and M. Padberg, “On the symmetric travelling salesman problem I: inequalities,” *Mathematical Programming*, vol. 16, pp. 265–280, 1979.
- [10] M. Grötschel and M. Padberg, “On the symmetric travelling salesman problem II: lifting theorems and facets,” *Mathematical Programming*, vol. 16, pp. 281–302, 1979.

- [11] M. Grötschel and W. Pulleyblank, “Clique tree inequalities and the symmetric travelling salesman problem,” *Mathematics of Operations Research*, vol. 11, pp. 537–569, 1986.
- [12] M. Jünger, G. Reinelt, and G. Rinaldi, “The traveling salesman problem,” in *Network Models* (M. Ball, T. Magnanti, C. Monma, and G. Nemhauser, eds.), vol. 7 of *Handbooks in Operations Research and Management Science*, ch. 4, pp. 225–330, Amsterdam: North-Holland, 1995.
- [13] E. Lawler, J. Lenstra, A. Rinnooy Kan, and D. Shmoys, eds., *The traveling salesman problem*. Wiley-Interscience Series in Discrete Mathematics, Chichester: John Wiley & Sons, 1985.
- [14] D. Naddef, “Polyhedral theory, branch and cut algorithms for the symmetric traveling salesman problem,” in *The Traveling Salesman Problem and its Variations* (G. Gutin and A. Punnen, eds.), Combinatorial Optimization, pp. 29–116, Dordrecht, NL: Kluwer Academic Publisher, 2002.
- [15] D. Naddef and Y. Pochet, “The symmetric traveling salesman polytope revisited,” *Mathematics of Operations Research*, vol. 26, pp. 700–722, 2001.
- [16] D. Naddef and G. Rinaldi, “The symmetric traveling salesman polytope: new facets from the graphical relaxation,” Tech. Rep. 248, IASI-CNR, 1988.
- [17] D. Naddef and G. Rinaldi, “The symmetric traveling salesman polytope and its graphical relaxation: composition of valid inequalities,” *Mathematical Programming*, vol. 51, pp. 359–400, 1991.
- [18] D. Naddef and G. Rinaldi, “The graphical relaxation: a new framework for the symmetric traveling salesman polytope,” *Mathematical Programming*, vol. 58, pp. 53–88, 1993.
- [19] D. Naddef and S. Thienel, “Efficient separation routines for the symmetric traveling salesman problem I: general tools and comb separation,” *Mathematical Programming*, vol. 92-2, pp. 237–255, 2002.
- [20] D. Naddef and S. Thienel, “Efficient separation routines for the symmetric traveling salesman problem II: separating multi handle inequalities,” *Mathematical Programming*, vol. 92-2, pp. 257–285, 2002.
- [21] D. Naddef and E. Wild, “The domino inequalities: facets for the symmetric traveling salesman polytope,” *Mathematical Programming*, vol. 98, pp. 223–251, 2003.
- [22] G. Nemhauser and L. Wolsey, *Integer and combinatorial optimization*. Discrete Mathematics and Optimization, New York, USA: Wiley-Interscience, 1988.
- [23] M. Oswald, G. Reinelt, and D. Theis, “Not every GTSP facet induces a STSP facet,” in *Integer Programming & Combinatorial Optimization (IPCO) XI* (M. Jünger and V. Kaibel, eds.), vol. 3509 of *Lecture Notes in Computer Science*, pp. 468–482, Springer-Verlag GmbH, 2005.
- [24] M. Padberg and S. Hong, “On the symmetric travelling salesman problem: a computational study,” *Mathematical Programming Study*, vol. 12, pp. 78–107, 1980.