

ISTITUTO DI ANALISI DEI SISTEMI ED INFORMATICA
“Antonio Ruberti”
CONSIGLIO NAZIONALE DELLE RICERCHE

**J. Li, M. Wang, A. De Gaetano, P. Palumbo,
S. Panunzi**

**SOME RESULTS ON THE GLOBAL STABILITY
OF THE EQUILIBRIUM FOR AN IVGTT MODEL**

R. 12, 2012

Jiaxu Li – Department of Mathematics, University of Louisville, Louisville, KY 40292.
jiaxu.li@louisville.edu.

Minghu Wang – Department of Mathematics, University of Louisville, Louisville, KY 40292.

Andrea De Gaetano – Istituto di Analisi dei Sistemi ed Informatica “A. Ruberti”, Consiglio Nazionale delle Ricerche (IASI-CNR), BioMatLab - UCSC - Largo A. Gemelli 8, 00168 Rome, Italy.

Pasquale Palumbo – Istituto di Analisi dei Sistemi ed Informatica “A. Ruberti”, Consiglio Nazionale delle Ricerche (IASI-CNR), BioMatLab - UCSC - Largo A. Gemelli 8, 00168 Rome, Italy. pasquale.palumbo@iasi.cnr.it.

Simona Panunzi – Istituto di Analisi dei Sistemi ed Informatica “A. Ruberti”, Consiglio Nazionale delle Ricerche (IASI-CNR), BioMatLab - UCSC - Largo A. Gemelli 8, 00168 Rome, Italy.

Collana dei Rapporti dell'Istituto di Analisi dei Sistemi ed Informatica "Antonio Ruberti",
CNR

viale Manzoni 30, 00185 ROMA, Italy

tel. ++39-06-77161

fax ++39-06-7716461

email: iasi@iasi.cnr.it

URL: <http://www.iasi.cnr.it>

Abstract

Diabetes mellitus has become a prevalent disease in the world. Diagnostic protocol for the onset of diabetes mellitus is the initial step in the treatments. The intravenous glucose tolerance test (IVGTT) has been considered as the most accurate method to determine the insulin sensitivity and glucose effectiveness. It is well known that there exists a time delay in insulin secretion stimulated by the elevated glucose concentration level. However, the range of the length of the delay in the existing IVGTT models are not fully discussed and thus in many cases the time delay may be assigned to a value out of its reasonable range. In addition, several attempts had been made to determine when the unique equilibrium point is globally stable. However, all these conditions are delay-independent. In this paper, we discuss the range of the time delay and provide easy-to-check delay-dependent conditions for the global stability of the equilibrium point for a recent IVGTT model through Liapunov function approach. Estimates of the upper bound of the delay for global stability are given in corollaries. In addition, the numerical simulation in this paper is fully incorporated with functional initial conditions, which is natural and more appropriate in delay differential equation system.

Key words: IVGTT; insulin secretion time delay; global stability; Liapunov function.

SOME RESULTS ON THE GLOBAL STABILITY OF THE EQUILIBRIUM FOR AN IVGTT MODEL

JIAXU LI*, MINGHU WANG[†], ANDREA DE GAETANO[‡], PASQUALE PALUMBO[‡], AND SIMONA PANUNZI[‡]

Abstract. Diabetes mellitus has become a prevalent disease in the world. Diagnostic protocol for the onset of diabetes mellitus is the initial step in the treatments. The intravenous glucose tolerance test (IVGTT) has been considered as the most accurate method to determine the insulin sensitivity and glucose effectiveness. It is well known that there exists a time delay in insulin secretion stimulated by the elevated glucose concentration level. However, the range of the length of the delay in the existing IVGTT models are not fully discussed and thus in many cases the time delay may be assigned to a value out of its reasonable range. In addition, several attempts had been made to determine when the unique equilibrium point is globally stable. However, all these conditions are delay-independent. In this paper, we discuss the range of the time delay and provide easy-to-check delay-dependent conditions for the global stability of the equilibrium point for a recent IVGTT model through Liapunov function approach. Estimates of the upper bound of the delay for global stability are given in corollaries. In addition, the numerical simulation in this paper is fully incorporated with functional initial conditions, which is natural and more appropriate in delay differential equation system.

Key words. IVGTT, insulin secretion time delay, global stability, Liapunov function

1. Introduction. Human bodies need to maintain a glucose concentration level in a narrow range (70-109 ml/dl or 3.9-6.04 mmol/l after overnight fast) ([10]). If one's glucose concentration level is significantly out of the normal range, this person is considered to have a plasma glucose problem: hyperglycemia or hypoglycemia. Diabetes mellitus, or simply, diabetes, is characterized by high blood glucose levels resulted from defects in the glucose-insulin endocrine metabolic regulatory system, in which either the pancreas does not release insulin, or the cells do not properly use insulin to uptake glucose.

Diabetes mellitus has become an epidemic disease in the sense of life style. To diagnose whether or not an individual subject is already a diabetic or has the potential to develop such disease, the so-called metabolic portrait, including the insulin sensitivity index and glucose effectiveness, of the subject needs to be sketched. To this end, several glucose tolerance tests have been developed and applied in clinics and experiments ([93], [10], [8], [41], [76], [16] and [61]). The fundamental idea of such tests is to examine the response of insulin, called insulin sensitivity, after a large amount of glucose is infused in to one's body. A commonly used protocol is the intravenous glucose tolerance test (IVGTT). In the procedure of IVGTT, overnight fast is required for the subject, and then the subject is given a bolus of glucose infusion intravenously, for example, 0.33 g/kg body weight [23] or 0.5 g/kg body weight of a 50% solution, and is administered into an antecubital vein in approximately 2 minutes. To observe the metabolic regulation between the glucose and insulin, within the next 180 minutes, the plasma glucose and serum insulin of the subject are sampled frequently at the time marks 2', 4', 6', 8', 10', 12', 14', 18', 21', 24', 30', 35', 40', 45', 50', 60', 70',

*Department of Mathematics, University of Louisville, Louisville, KY 40292. Jiaxu Li is supported in part by NIH Grant R01-DE019243, P30ES01443 and DOE Grant DE-EM0000197. (jiaxu.li@louisville.edu)

[†]Department of Mathematics, University of Louisville, Louisville, KY 40292. Minghu Wang is supported by DOE DE-EM0000197.

[‡]CNR-IASI BioMatLab, Largo A. Gemelli 8 - 00168 Rome, Italy.

80', 90', 100', 120', 140', 160' and 180'. According to the rich information constituted in the sampled data, appropriate analysis can reveal the metabolic portrait.

One popular approach of the analysis is as follows: 1) formulate or choose a well-formulated kinetic model based on physiology; 2) estimate the parameters of the IVGTT model with experimental data, and then 3) the parameter values are used to obtain physiological information, for example, the insulin sensitivity.

Several IVGTT models have been proposed and some are widely used ([2], [4], [9], [11], [14], [16], [23]). All these models incorporate the insulin secretion delay implicitly in ODE model by using compartment-split technique or explicitly in DDE models. However, none of above work has discussed the reasonable range for the time delay and how to estimate the reasonable value of the delay according to the sampled data. Instead, the length of delay is often set to be larger than 20 minutes, which is in contrast to that the delay is between 5 and 15 minutes for normal subjects in normal environment ([17], [10], [22]). Intuitively, the length of the delay in glucose tolerance test should not be longer than that in normal environment. In addition, several attempts have been made to obtain the conditions for the global stability of the equilibrium point of the models ([6], [9], [14] and [16]). Unfortunately all the conditions are delay-independent. For the information about the delay $\tau > 0$, it is only known that there exists a $\tau_0 > 0$ such that the equilibrium point is locally stable if $\tau < \tau_0$; and unstable otherwise (thus a periodic solution is bifurcated out). To reveal the insight regarding the time delay, it would be very helpful to know an estimate of the upper bound $\bar{\tau}$ so that the equilibrium point is globally stable when $\tau \leq \bar{\tau}$. Applying the Liapunov function approach, we obtain delay dependent conditions for the global stability of the single delay IVGTT model proposed in [14]. Estimated upper bounds can be derived from these conditions explicitly or implicitly. We also improve the numerical simulations by using functional initial conditions in the delay differential equation model. To simulate the bolus glucose infusion and the abrupt insulin secretion, constant initial conditions with adjustments at time $t = 0$ are applied in [4], [6], [9], [11], [14], [16]. This is neither natural to the design of the test, nor appropriate for a functional differential equation system. In this paper, we will shift the starting time of glucose infusion to -2' and use a non-constant function as initial value in $[-\tau, 0]$.

We organize this paper as follows. We will first discuss the range of the length of delay, then obtain delay-dependent conditions for the global stability of the equilibrium point, from which estimated upper bounds of τ_0 are given in corollaries, and finally show numerical simulations for DDE IVGTT models. We first in Section 3 introduce the IVGTT models proposed and studied in [6], [14] and [16], followed by the analysis of the models, in which delay dependent conditions of global stability of the unique equilibrium are given by utilizing Liapunov function in Section 4. In Section 5 we perform numerical simulations and we end this paper with discussions in Section 6.

2. Range of the time delay of insulin secretion. All discrete and distributed delay differential equation models for IVGTT involve explicitly a time delay between the rise in glycemia and the correspondingly stimulated insulin secretion. Thus, it is intuitive to ask questions: What is the definition of the time delay? How long is the delay? De Gaetano and Arino ([4]) considered as indicative of the overall length of delay the average of delays of all studied subjects. All IVGTT models in [6], [9], [11], [14] and [16] follow the same criterion, which yields a value of around 20 minutes.

The events of the insulin secretion from pancreatic β -cells can be summarized as

follows according to [10] and the references therein. As the plasma glucose concentration is elevated, the insulin secretion from β -cells is determined by several cascading complex electric processes inside islet. These processes can be described in following steps: a glucose transporter GLUT2 transports glucose molecules into islet, the ratio of ATP (adenosine triphosphate) over ADP (adenosine diphosphate) is increased, the K^+ channels are closed and consequently the Ca^{2+} channels open, and then the influx of Ca^{2+} ions triggers insulin exocytosis from β -cell granules. Obviously such a chain of events causes a time delay for insulin release in responding to elevated glucose concentrations. Insulin release has effects on both hepatic glucose production and insulin-dependent glucose utilization ([3]). It takes some time for the newly synthesized insulin to cross the endothelial barrier and eventually become the so-called “remote insulin”, then act on glucose uptake. This total delay time is approximately 5 to 15 min ([3], [22], [24]).

Bolus glucose infusion causes abrupt increase in plasma glucose concentration. Such an abrupt glucose increase stimulates bi-phasic insulin secretions, a rapid and approximately linear but short first phase release, followed by a slower and longer second phasic secretion. According to Overgaard et al. ([13]) and the references therein, such secretions relate to multiple aspects including the impact of glucose on β -cells, insulin synthesis, movements of insulin in β -cells, and eventually insulin release from granules. As glucose is infused intravenously, the metabolic system responds quickly. The rate of proinsulin synthesis is approximately linearly increased in translation of proinsulin mRNA. The proinsulin is split into insulin and C-peptide and stored in granules. A fraction of such granules is docked on the surface of the plasma membrane while others stay in the intracellular space with the ability of free movement. The group of granules docked on the surface of the plasma membrane is called readily-releasable pool (RRP), the insulin in which is released at the quick infusion of the glucose and forms the first phase of secretion. This release is instantaneous or almost instantaneous since it takes almost no time for the glucose infuse in intravenous to transport to pancreas and the insulin in the ready releasable pool. This instantaneous insulin release is followed by the following actions: grandules in intracellular space flow into the RRP and then their insulin is released. This release forms the second phase of secretion. This series of chemical and electrical events causes a time lag in the release after the observation of the increased glucose concentration. This time lag can be a few minutes ([13]).

Apparently, second phase secretion in IVGTT is not significantly distinguishable from the secretion under normal conditions (meals). Therefore time delay should exist and the range of the delay should not be longer than that in normal status. According to physiology cited in their work, Li et al. ([10]) defined the delay as the length of the delay τ as “from the time that the glucose concentration level is elevated to the moment that the insulin has been transported to interstitial space and becomes remote insulin”, and determined that the possible value of the delay falls in the biological range of 5 minutes to 15 minutes ([17]). Therefore we suspect that for most subjects, the length of the delay is shorter than 15 minutes in IVGTT. In most of the previous work, the time delay was chosen between 18 minute and 24 minutes. The mean value, according to [14] and [16], is 19.271 minutes. It must be underscored that no direct relationship between the estimated parameter value and morphological features of the resulting state variable timecourse can be drawn, even though, empirically, the delay often approximates the time at which the trough between primary and secondary insulin releases is reached.

It is well known that a large delay can destabilize a system ([7]). An accurate assessment of the delay can therefore play a critical role in the establishment of the stability properties of the model.

3. Single delay models for intravenous glucose tolerance test.

3.1. The models. The Minimal Model, proposed by Bergman and his colleagues in 1979 ([2]) and 1980 ([23]), is believed to be one of the first widely employed models for the IVGTT ([12]). The Minimal Model is in fact a combination of two separate ordinary differential equation (ODE) models, one for glucose kinetics and one for insulin kinetics. The time delay in the glucose-insulin regulatory system was simulated by the chain trick through an auxiliary variable ([21]). However, certain mathematical problems arise in this unified Minimal Model, for example, the model fails to exhibit any equilibrium point and one of the state variables (insulin activity at the distant site) increases with time without any upper bound. In 2000, De Gaetano and Arino ([4]) pointed out these issues and proposed the first delay differential equation (DDE) model, called *dynamic model*. This novel DDE model is not only well-posed in mathematics, but also more physiologically appealing as the time delay is contained in the model explicitly. The dynamic model was then generalized by Li et al. ([9]) in 2001. Another specific delay model proposed in 2004 by Mukhopadhyay et al. ([11]) also falls within the wide class of models contemplated by Li et al. ([9]). Finally, De Gaetano and colleagues proposed a specific functional form of a single discrete delay model (SDM), motivated its parameter choice physiologically, estimated it comparatively with the Minimal Model ([16]) showing greater robustness and precision, and proved several of its satisfactory qualitative properties ([14]). Finally, the same authors extended the class of subjects tested to cover a wide spectrum of insulin resistance and pointed to the drawbacks of "decoupling" estimation methods as typically employed for the Minimal Model (advantages of the single delay model for the assessment of insulin sensitivity from the intravenous glucose tolerance test, [15]).

Denote the plasma concentrations of glucose and insulin at time $t \geq -\tau$ by $G(t) > 0$ (mg/dl) and $I(t) > 0$ (μ U/ml), respectively. The SDM model is given by ([16]).

$$(3.1) \quad \begin{cases} G'(t) = b - eG(t) - aG(t)I(t) \\ I'(t) = df(G(t-\tau)) - cI(t) \end{cases}$$

with initial conditions $G(t) \equiv G_b$ for $t \in [-\tau, 0)$, $G(0) = G_b + G_\Delta$ and $I(0) = I_b + I_\Delta G_\Delta$, where $G_b > 0$ and $I_b > 0$ basal levels of glycemia and insulinemia after overnight fast (also called baseline), and G_Δ and I_Δ are the theoretical increase over basal concentrations. The parameter $b > 0$ (mg/dl/min) is the constant glucose input; $e \geq 0$ (1/min) is the insulin independent glucose utilization by, e.g., brain cells; $a > 0$ (ml/ μ U/min) is the insulin dependent glucose utilization; $c > 0$ (1/min) is the insulin degradation; and the term $df(G(t-\tau))$ is the insulin secretion responded to the glucose stimulation with time delay $\tau \geq 0$

$$f(x) = \frac{x^\gamma}{\alpha^\gamma + x^\gamma}$$

with $d > 0$ (μ U/mg/min) as the maximum secretion, $\alpha > 0$ as the half-saturation and $\gamma > 0$. If $\gamma \leq 1$, $f(x)$ is in the shape of Michaelis-Menten kinetics; if $\gamma > 1$, $f(x)$

is in sigmoidal shape. As discussed in [14], the model 3.1 has a unique equilibrium and the equilibrium point is at (G_b, I_b) , so that after a perturbation the glucose and insulin blood levels will return back their basal value, which implies that

$$(3.2) \quad b = G_b(e + aI_b) \quad \text{and} \quad d = cI_b/(1 + (\alpha/G_b)^\gamma).$$

It must be noticed that the SDM also falls in the general class of models studied by Li et al. in [9]. Therefore what analytical results were obtained in [9] automatically apply for the SDM. In the present work, in order to simplify mathematical notation at the possible expense of physiological clarity, we use a different notation for the parameters with respect to (3.1) from those in [6], [14] and [16]. The relations are as follow: $a = K_{xgi}$, $b = T_{gh}/V_G$, $c = K_{xi}$, $d = T_{iG_{max}}/V_I$, and $e = K_{xg}$. According to both the Minimal Model ([2], [23]) and the SDM (3.1) ([16]), the parameter a is called insulin sensitivity index, and the parameter e is called glucose effectiveness.

The main determinants of glucose uptake from plasma in the body are brain glucose consumption (essentially constant, therefor represented as a zero-order kinetics) and muscle and adipose tissue consumption and storage (insulin and glucose-concentration dependent, hence second-order kinetics). Therefore, first-order, insulin-independent glucose utilization has often been considered negligible([6], [14] and [16]), i.e., $e = 0$. Therefore model (3.1) can be further simplified as follows ([16])

$$(3.3) \quad \begin{cases} G'(t) = b - aG(t)I(t) \\ I'(t) = df(G(t - \tau)) - cI(t) \end{cases}$$

The authors of [16] applied the Akaike Information Criterion (AIC) in model selection by comparing model (3.3) with model (3.1), and determined that model (3.3) is more suitable for representing observed IVGTTs. Moreover, impaired insulin-dependent glucose disposal is one of the reasons of insulin resistance that is typical in type 2 diabetes. Decreased insulin-dependent glucose transport by GLUT4 also occurs, for example, in chronic alcohol users ([18]). Low insulin sensitivity may also occur in type 1 diabetes and could be caused by impaired GLUT4 translocation ([5]). This indicates that insulin-independent GLUT1 may be dominant in translocation of glucose molecules ([5]) at least in some disease states. These considerations, as well as potential concentration effects shifting (at least in the short term) glucose to the interstitial space, prompt a cautious re-evaluation of the role of insulin-independent tissue glucose uptake and therefore of the generally positive, if small, value of the parameter e .

3.2. Existing analytical results and numerical observations. While analysis of the original Minimal Model focused only on numerical simulations ([2] and [23]), a number of analytical studies for subsequent IVGTT models have been carried out ([4], [6], [9], [11], [14]). Typically, these last authors studied the local stability of the unique steady state and then numerically demonstrated that larger delays would not destabilize the steady state, since numerically determined Hopf bifurcation points are far beyond the meaningful physiological range. Some analytical delay-independent conditions for global stability of the steady state were obtained for the models proposed by [6], [9], and [14]. Particularly, when $\gamma \leq 1$, the unique equilibrium point is globally asymptotically stable by applying the Theorem 3.1 in [9] with simple computations, and a set of sufficient and necessary conditions is given for the model (3.1) ([14]). However, physiology ([8], [10], [12], [17], [19], [22], [24]) and all the clinical

data ([6], [14], [16]) show the existence of the time delay of the glucose stimulated insulin secretion. Existing delay dependent condition (Theorem 3 in [6]) on the global stability is not satisfied with clinical data (Remark 11 in [14]), although the conditions hold with some critical modification, for example, the time delay has to be smaller (Remark 9 in [6]). Nevertheless, the local stability of the equilibrium and numerically determined large Hopf bifurcation value of the time delay do not destabilize the equilibrium point for a wide reasonable set of parameter values ([8], [14]). Studies and applications of these models are extended by Panunzi et al. ([16]) in application, and Giang et al. ([6]) in analysis. For more details for the formulation and applications of model (3.1) and model (3.3), interested readers can refer to [6], [14], and [16].

We summarize the analytical results obtained in above references in the following. The system (3.1) is called persistent if the solutions are eventually uniformly bounded by positive constants from both above and below. A fastly oscillated solution is defined as a solution $(G(t), I(t))$ that oscillates around the equilibrium (G_b, I_b) and within two consecutive zeros $t_0 < t_1$ of $I(t) - I_b$, $I(t)$ attains its maximum or minimum at $t_* < t_0 + 2\tau$. (Definition 6 and Theorem 3 in [6]).

Let $(G(t), I(t))$ be a solution of (3.1) or model (3.3). Throughout this paper, we define

$$\bar{G} = \limsup_{t \rightarrow \infty} G(t), \quad \underline{G} = \liminf_{t \rightarrow \infty} G(t) \quad \bar{I} = \limsup_{t \rightarrow \infty} I(t), \quad \underline{I} = \liminf_{t \rightarrow \infty} I(t).$$

- (A1) The model (3.1) and the model (3.3) are persistent (Proposition 2.2 in [9]; Theorem 2 in [14]). Thus, the limits defined above are finite.
 (A2) Following inequalities hold (Lemma 2.1 in [9]; Remark 5 in [14]).

$$(3.4) \quad df(\underline{G}) \leq c\underline{I} \leq c\bar{I} \leq df(\bar{G})$$

$$(3.5) \quad (e + a\underline{I})\bar{G} \leq b \leq (e + a\bar{I})\underline{G}$$

- (A3) The necessary condition (Theorem 5.1 in [9]; Theorem 6 in [14]) and sufficient conditions (Theorem 4 in [14]) for global stability of model (3.1) and model (3.3) is

$$(3.6) \quad \frac{\gamma a I_b}{1 + \left(\frac{\underline{G}_b}{\alpha}\right)^\gamma} \leq e + a I_b.$$

- (A4) If $\gamma \leq 1$, the unique equilibrium the model (3.1) or the model (3.3) is globally asymptotically stable (apply Theorem 3.1 in [9] to this special case). Notice that this result does not need the assumption (3.2) or (3.6). So we only consider the case $\gamma > 1$ in this paper.
 (A5) If the inequality in (3.6) does not hold, then there is a $\tau_0 > 0$ such that (i) when $\tau < \tau_0$, the equilibrium (G_b, I_b) is locally stable; (ii) when $\tau \geq \tau_0$, the equilibrium is unstable. In other words, the model (3.1) and the model (3.3) undergo a Hopf bifurcation at $\tau = \tau_0$ when τ increases from $\tau = 0$ (Theorem 6 in [14]).
 (A6) In the model (3.1), the equilibrium is a global attractor if

$$\frac{adG_b}{ce} \sqrt{\sup_{[0, G_b]} f'(G) \sup_{[G_b, \infty)} f'(G)} < 1.$$

(Theorem 2 in [6].)

(A7) Every fastly oscillated solution of the model (3.3) converges to the positive equilibrium if

$$(1 - e^{-2c\tau}) \frac{adG_b}{ceI_b} \sqrt{G_b M_* \sup_{[0, G_b]} f'(G) \sup_{[G_b, \infty)} f'(G)} < 1,$$

where M_* satisfies $m_* f(M_*) = M_* f(m_*) = G_b f(G_b)$ for $m_* < G_b < M_*$. (Theorem 3 in [6].)

Notice that the conditions for the globally asymptotical stability of the equilibrium in (A3), (A4) and (A6) are delay-independent. But (A3) is a necessary and sufficient condition, so the result in (A5) shall be either a weaker condition or equivalent to (A3). Nevertheless, according to [14], only 3% of clinical data satisfy the condition in (A3). So to ensure the global attractive behavior of the tolerance test, analysis on delay dependent global stability is necessary. Notice that the insulin dynamics in IVGTT may often reach its basal level without oscillation ([14], [16]) as defined in [6], the application of the result in (A6) is limited. In addition, (A6) requires τ to be small enough so that the parameters fitted from the clinical data satisfy the condition ([6], Remark 9.)

4. Global stability of the equilibrium. We will establish criteria to determine the global stability, which depend on the time delay, and thus provide the estimated upper bounds for time delay.

We state following well known fact without proof, which is useful in estimating the bounds of a solution $(G(t), I(t))$ of the model (3.1).

LEMMA 4.1. *If $h'(t) \leq p - qh(t)$, or $h'(t) \geq p - qh(t)$, where $p, q > 0$, then*

$$h(t) \leq \frac{p}{q} + \left(h(0) - \frac{p}{q}\right)e^{-qt}, \quad \text{or} \quad h(t) \geq \frac{p}{q} + \left(h(0) - \frac{p}{q}\right)e^{-qt},$$

for all $t \geq 0$.

A direct application of Lemma 4.1 is to estimate the bounds of a solution $(G(t), I(t))$.

We have

LEMMA 4.2. *If $(G(t), I(t))$ is a solution with positive initial conditions of the model (3.1), then*

$$m_I \doteq \min \left\{ I(0), \frac{k_2}{c} \right\} \leq I(t) \leq M_I \doteq \max \left\{ I(0), \frac{d}{c} \right\}$$

and

$$m_G \doteq \min \left\{ G(0), \frac{b}{k_1} \right\} \leq G(t) \leq M_G \doteq \max \left\{ G(0), \frac{b}{k_3} \right\}.$$

where $k_1 = e + aM_I$, $k_2 = df(m_G)$ and $k_3 = e + am_I$.

The proof is straight forward in observing the positiveness of such solutions (Theorem 1 in [14]).

Remark The bounds given by the lemma 4.2 is more accurate than the consequence in A2 by [14], which would result in

$$\frac{bc}{ec + ad} \leq \underline{G} \leq \overline{G} \leq \frac{bc}{ec + adf\left(\frac{bc}{ec+ad}\right)}.$$

by simple computation. However the upper bound M_G of $G(t)$ for $t > 0$ rather than the bound at limiting status of $G(t)$ plays an important role when applying the theorem 4.3 to determine the global stability of the equilibrium. Notice that M_G is dependent of the initial values $G(0)$ and $I(0)$ of the solution $(G(t), I(t))$ only.

For the sake of convenience, we denote for $\gamma > 1$

$$(4.1) \quad L = d \frac{(\gamma + 1)^2}{4\gamma\alpha} \left(\frac{\gamma - 1}{\gamma + 1} \right)^{\frac{\gamma-1}{\gamma}},$$

which is the maximum value of the derivative of $df(G)$ for $G > 0$, and

$$(4.2) \quad R = e + aI_b, \quad \text{and} \quad K = aM_G.$$

By Lemma 4.2, L, R and K are independent of τ . When consider the model (3.3), $R = aI_b$.

When $\tau = 0$, model (3.1) and (3.3) become a two-dimensional ordinary differential equation system. It is easy to show that the unique equilibrium (G_b, I_b) is globally asymptotically stable by Poincare-Bendixson Theorem. When $\tau > 0$, the models are functional differential equations. Poincare-Bendixson Theorem does not apply. To establish a criterion of delay dependent global stability, we build a Liapunov function and prove that the solutions of model (3.1) or model (3.3) are attracted to the equilibrium (G_b, I_b) .

THEOREM 4.3. *If there exists a positive constant $u > 0$ such that*

$$(4.3) \quad \tau < \min \left\{ \frac{2u}{L}, \frac{2c}{L(R + 2K)} \right\},$$

and

$$(4.4) \quad \Delta \doteq M_u^2 - 4R \left(u - \frac{\tau}{2} L \right) \left(c - \frac{\tau}{2} L(R + 2K) \right) < 0,$$

where $M_u = \max\{L, Ku\}$, then the equilibrium (G_b, I_b) of model (3.1) is globally and asymptotically stable.

Proof. Let

$$(4.5) \quad W(G(t), I(t)) = \frac{1}{2}u(G(t) - G_b)^2 + \frac{1}{2}(I(t) - I_b)^2,$$

$$(4.6) \quad U(G(t), I(t)) = C \int_{t-\tau}^t \int_z^t (G(s) - G_b)^2 ds dz + D \int_{t-\tau}^t \int_z^t (I(s) - I_b)^2 ds dz.$$

where $C = \frac{1}{2}RL$ and $D = \frac{1}{2}KL$.

Let

$$V(G(t), I(t)) = W(G(t), I(t)) + U(G(t), I(t)),$$

then, clearly, V is a Liapunov function.

By Mean Value Theorem, we have

$$\begin{aligned} d(f(G(t - \tau)) - f(G_b)) &= df'(\xi)(G(t - \tau) - G_b) \\ &= df'(\xi) \left((G(t - \tau) - G(t)) + (G(t) - G_b) \right) \end{aligned}$$

where ξ is between $G(t - \tau)$ and $G(t)$. Also, since $uv \leq \frac{1}{2}(u^2 + v^2)$ for all $u, v \geq 0$, we have

$$\begin{aligned}
& |(I(t) - I_b)(G(t) - G(t - \tau))| = \left| (I(t) - I_b) \int_{t-\tau}^t G'(s) ds \right| \\
&= \left| \int_{t-\tau}^t (b - eG(s) - aG(s)I(s)) (I(t) - I_b) ds \right| \\
&= \left| \int_{t-\tau}^t (e(G(s) - G_b)(I(t) - I_b) + aG(s)(I(s) - I_b)(I(t) - I_b) \right. \\
&\quad \left. + aI_b(G(s) - G_b)(I(t) - I_b)) ds \right| \\
&= \left| \int_{t-\tau}^t ((e + aI_b)(G(s) - G_b)(I(t) - I_b) + aG(s)(I(s) - I_b)(I(t) - I_b)) ds \right| \\
&\leq \frac{1}{2} \left[(e + aI_b) \left(\int_{t-\tau}^t (G(s) - G_b)^2 ds + \tau(I(t) - I_b)^2 \right) \right. \\
&\quad \left. + aG_M \left(\int_{t-\tau}^t (I(s) - I_b)^2 ds + \tau(I(t) - I_b)^2 \right) \right] \\
&\leq \frac{1}{2} \left[R \int_{t-\tau}^t (G(s) - G_b)^2 ds + K \int_{t-\tau}^t (I(s) - I_b)^2 ds \right. \\
&\quad \left. + \tau(R + K)(I(t) - I_b)^2 \right].
\end{aligned}$$

Then

$$\begin{aligned}
\frac{d}{dt}W &= u(G(t) - G_b) \left[-e(G(t) - G_b) - a(G(t)(I(t) - I_b) + I_b(G(t) - G_b)) \right] \\
&\quad + (I(t) - I_b) \left[d(f(G(t - \tau)) - f(G_b)) - c(I(t) - I_b) \right] \\
&= -ue(G - G_b)^2 - uaG(G - G_b)(I - I_b) - uaI_b(G - G_b)^2 \\
&\quad + d(I - I_b) \left[f(G(t - \tau)) - f(G_b) \right] - c(I - I_b)^2 \\
&= -(ue + uaI_b)(G - G_b)^2 - uaG(G - G_b)(I - I_b) - c(I - I_b)^2 \\
&\quad + d(I - I_b) \left[f(G(t - \tau)) - f(G_b) \right] \\
&= -u(e + aI_b)(G - G_b)^2 - uaG(G - G_b)(I - I_b) - c(I - I_b)^2 \\
&\quad + df'(\xi)(I - I_b) \left[(G(t - \tau) - G(t)) + (G(t) - G_b) \right] \\
&= -u(e + aI_b)(G - G_b)^2 + (df'(\xi) - uaG)(G - G_b)(I - I_b) - c(I - I_b)^2 \\
&\quad + df'(\xi)(I - I_b)(G(t - \tau) - G(t)) \\
&\leq -u(e + aI_b)(G - G_b)^2 + |df'(\xi) - uaG||G - G_b||I - I_b| - c(I - I_b)^2 \\
&\quad + L|(I - I_b)(G(t - \tau) - G(t))| \\
&\leq -uR(G - G_b)^2 + |df'(\xi) - uaG||G - G_b||I - I_b| - c(I - I_b)^2 \\
&\quad + \frac{1}{2}L \left[R \int_{t-\tau}^t (G(s) - G_b)^2 ds + K \int_{t-\tau}^t (I(s) - I_b)^2 ds \right. \\
&\quad \left. + \tau(R + K)(I(t) - I_b)^2 \right].
\end{aligned}$$

$$\begin{aligned}
&= -uR(G - G_b)^2 + |df'(\xi) - uaG||G - G_b||I - I_b| \\
&\quad - \left(c - \frac{\tau}{2}L(R + K)\right)(I - I_b)^2 \\
&\quad + C \int_{t-\tau}^t (G(s) - G_b)^2 ds + D \int_{t-\tau}^t (I(s) - I_b)^2 ds
\end{aligned}$$

Furthermore,

$$\begin{aligned}
\frac{dU}{dt} &= -C \int_{t-\tau}^t (G(s) - G_b)^2 ds + C\tau(G(t) - G_b)^2 \\
&\quad - D \int_{t-\tau}^t (I(s) - I_b)^2 ds + D\tau(I(t) - I_b)^2
\end{aligned}$$

So, along the solution $(G(t), I(t))$ of model (3.1), we have

$$\begin{aligned}
\frac{dV}{dt} &\leq -uR(G - G_b)^2 + |df'(\xi) - uaG||G - G_b||I - I_b| \\
&\quad - \left(c - \frac{\tau}{2}L(R + K)\right)(I - I_b)^2 \\
&\quad + C \int_{t-\tau}^t (G(s) - G_b)^2 ds + D \int_{t-\tau}^t (I(s) - I_b)^2 ds \\
&\quad - C \int_{t-\tau}^t (G(s) - G_b)^2 ds + C\tau(G(t) - G_b)^2 \\
&\quad - D \int_{t-\tau}^t (I(s) - I_b)^2 ds + D\tau(I(t) - I_b)^2 \\
&= -(uR - C\tau)(G - G_b)^2 + |df'(\xi) - uaG||G - G_b||I - I_b| \\
&\quad - \left(c - \tau\left(\frac{1}{2}L(R + K) + D\right)\right)(I - I_b)^2 \\
&= -\left(uR - \frac{\tau}{2}LR\right)(G - G_b)^2 + |df'(\xi) - uaG||G - G_b||I - I_b| \\
&\quad - \left(c - \tau\left(\frac{1}{2}L(R + K) + \frac{1}{2}LK\right)\right)(I - I_b)^2 \\
&= -R\left(u - \frac{\tau}{2}L\right)(G - G_b)^2 + |df'(\xi) - uaG||G - G_b||I - I_b| \\
&\quad - \left(c - \frac{\tau}{2}L(R + 2K)\right)(I - I_b)^2 \\
&\leq -R\left(u - \frac{\tau}{2}L\right)(G - G_b)^2 + M_u|G - G_b||I - I_b| \\
&\quad - \left(c - \frac{\tau}{2}L(R + 2K)\right)(I - I_b)^2 \\
&< 0
\end{aligned}$$

in observing that the conditions (4.3) and (4.4) hold. \square

Remark Since M_u can be dependent of $G(0)$ or $I(0)$ in some cases, the global stability is not uniform.

We derive two corollaries from the above theorem, which give the estimates of the bounds for the delay τ , which ensures the equilibrium is globally stable. Denote

$$p_1 = \frac{cK + L(R + 2K)}{KL(R + 2K)}, \quad q_1 = \frac{4Rc - KL}{KRL(R + 2K)}$$

and

$$\tilde{\tau} = \min \left\{ \frac{2}{K}, \frac{2c}{L(R+2K)} \right\}.$$

Then we have

COROLLARY 4.4. *The unique equilibrium of the model (3.1) is globally asymptotically stable, if the delay τ satisfies either of the following conditions:*

$$(4.7) \quad (i) \tau < \tilde{\tau}, \text{ if } p_1^2 - q_1 < 0;$$

$$(4.8) \quad (ii) \tau < \min\{\tilde{\tau}, p_1 - \sqrt{p_1^2 - q_1}\}, \text{ if } p_1^2 - q_1 > 0 \text{ and } q_1 > 0;$$

$$(4.9) \quad (iii) \tau \in (p_1 + \sqrt{p_1^2 - q_1}, \tilde{\tau}), \text{ if } p_1^2 - q_1 > 0 \text{ and } p_1 + \sqrt{p_1^2 - q_1} < \tilde{\tau};$$

$$(4.10) \quad (iv) \tau < \tilde{\tau} \text{ and } \tau \neq p_1, \text{ if } p_1^2 - q_1 = 0;$$

Proof. Notice that $M_u = \max\{L, Ku\}$ in Theorem 4.3. Let $u = L/K$, then $M_u = L$. Therefore (4.3) becomes $\tau < \tilde{\tau}$, and (4.4) holds if and only if

$$\begin{aligned} & M_u^2 - 4R\left(u - \frac{\tau}{2}L\right)\left(c - \frac{\tau}{2}L(R+2K)\right) \\ &= L^2 - 4R\left(\frac{L}{K} - \frac{\tau}{2}L\right)\left(c - \frac{\tau}{2}L(R+2K)\right) \\ &= L\left[L - 4R\left(\frac{1}{K} - \frac{\tau}{2}\right)\left(c - \frac{\tau}{2}L(R+2K)\right)\right] \\ &= L\left[L - 4R\left(\frac{c}{K} - \frac{\tau}{2}c - \frac{\tau}{2}\frac{L(R+2K)}{K} + \frac{\tau^2}{4}L(R+2K)\right)\right] \\ &= L\left[L - \frac{4Rc}{K} + 2\tau R\left(c + \frac{L(R+2K)}{K}\right) - \tau^2 RL(R+2K)\right] \\ &= -L^2R(R+2K)\left[\frac{-L + \frac{4Rc}{K}}{RL(R+2L)} - \frac{2R(c + \frac{L(R+2K)}{K})}{RL(R+2K)}\tau + \tau^2\right] \\ &= -L^2R(R+2K)\left[\frac{4Rc - KL}{KRL(R+2K)} - \frac{2(cK + L(R+2K))}{KL(R+2K)}\tau + \tau^2\right] \\ &= -L^2R(R+2K)(q_1 - 2p_1\tau + \tau^2) \\ &< 0. \end{aligned}$$

Then it is straight forward to obtain the conclusion. \square

Denote

$$H = c - \frac{\tau}{2}L(R+2K) \quad \text{and} \quad \bar{\tau} = \frac{2Rc}{L(R+K)^2}.$$

If $\tau < \bar{\tau}$, then

$$\begin{aligned} 2RH - K^2L\tau &= 2RH - K^2L\tau \\ &= 2R\left(c - \frac{\tau}{2}L(R+2K)\right) - K^2L\tau \\ &= 2Rc - (RL(R+2K) + K^2L)\tau \\ &= 2Rc - L(R(R+2K) + K^2)\tau \\ &= 2Rc - L(R^2 + 2RK + K^2)\tau \\ &= 2Rc - L(R+K)^2\tau \\ &> 0. \end{aligned}$$

Hence, we can define

$$u_0 = \frac{4RH + \sqrt{8RH(2RH - K^2L\tau)}}{2K^2} > 0.$$

Therefore, we have

COROLLARY 4.5. *The unique equilibrium of the model (3.1) is globally asymptotically stable if*

$$(4.11) \quad \tau < \min \left\{ \frac{2u_0}{L}, \bar{\tau} \right\}, \quad \text{and} \quad u_0 > \frac{L}{K}.$$

Proof. We choose an appropriate $u > 0$ so that the inequality (4.4) holds. Let $u > L/K$. Then $M_u = Ku$ in Theorem 4.3 and thus (4.4) holds if and only if

$$K^2u^2 - 4R(u - \frac{\tau}{2}L)H = K^2u^2 - 4RHu + 2RLH\tau < 0.$$

So the existence of such $u > 0$ is equivalent to

$$(4RH)^2 - 4K^2 \cdot 2RLH\tau = 8RH(2RH - K^2L\tau) > 0,$$

and the larger root u_0 of $P(u) = K^2u^2 - 4RHu + 2RKH\tau = 0$ is greater than L/K . Notice (4.11). We choose any $u \in (L/K, u_0)$ so that both conditions in the Theorem 4.3 are satisfied. \square

Remark The inequality of (i), (ii) and (iv) in corollary 4.4 provide estimated upper bounds of the time delay of insulin secretion stimulated by glucose. Since H in (4.11) is dependent of τ , one can estimate an upper bound of τ implicitly.

We will apply Corollary 4.4 and Corollary 4.5 in next section with experimental data obtained from [4] and [16].

5. Validation and numerical simulations. In order to exemplify the computation of the proposed stability criteria, we considered sets of parameter values consistent with adaptation to data from actual IVGTT experiments. Simulations are performed by using Matlab delay differential equation solver dde23 ([20]). We considered three sets of experimental data from [4] listed in Table 5.1, and two sets from [16] listed in Table 5.2. In the original datasets, the time mark 0' is the starting time of the bolus glucose infusion, and the time mark 2' is the first blood sampling time. For simplicity in expressing the functional initial condition in the Glycemia state variable, we introduce a -2' shift of the time marks (3.1).

The initial condition for a delay differential equation with maximum delay τ is a function defined in the interval $[-\tau, 0]$. We further depart from published models in hypothesizing that glycemia increase between the start of infusion to the first observed glycemia point follows an ascending linear ramp. Therefore we use a piece-wise linear function

$$(5.1) \quad \phi(t) = \begin{cases} G_b, & \text{for } t \in [-\tau, -2] \\ G_0 + \frac{G_0 - G_b}{2}t, & \text{for } t \in (-2, 0]; \end{cases}$$

while $I(0) = I_0$. Here G_b and I_b are taken from the first row at time mark -2' in the data tables and G_0 and I_0 are taken from the second row at time mark 0' from

TABLE 5.1

Experimental data published in [4]. The first column is the time in minute to sample the blood with a two-minute shift. The second and third columns are the data for subject 6. The first valley is at about 8' mark. The fourth and fifth columns are the data from for the subject 7 and its first valley is at about 12' mark. The sixth and seventh columns are the data for subject 8 and there is no clear valley.

min	G(mg/dl)	I(μ U/l)	G(mg/dl)	I(μ U/l)	G(mg/dl)	I(μ U/l)
-2	87.7358	67.9245	87.2117	38.5744	77.9874	57.9000
0	225.4717	413.2075	299.3711	179.4549	226.4151	1031.4000
2	214.1509	410.3774	259.9581	103.9832	228.9308	915.7000
4	203.7736	305.6604	253.2495	99.7904	203.7736	759.7000
6	200.0000	286.7925	244.0252	93.9203	201.2579	772.3000
8	195.2830	234.9057	225.5765	104.8218	196.2264	646.5000
10	192.4528	317.9245	223.8994	77.1488	183.6478	669.2000
13	174.5283	278.3019	203.7736	88.8889	173.5849	513.2000
18	158.4906	238.6792	188.6792	95.5975	148.4277	508.2000
23	150.0000	250.0000	170.2306	79.6646	123.2704	440.3000
28	131.1321	233.9623	150.9434	97.2746	115.7233	327.0000
33	118.8679	203.7736	134.1719	86.3732	100.6289	286.8000
38	115.0943	153.7736	119.9161	108.1761	95.5975	226.4000
48	106.6038	169.8113	101.4675	44.4444	85.5346	166.0000
58	93.3962	115.0943	89.7275	24.3187	75.4717	148.4000
78	82.0755	111.3208	85.5346	33.5430	72.9560	118.2000
98	77.3585	53.7736	85.5346	29.3501		
118	83.0189	46.2264	88.0503	37.7358	77.9874	67.9000
138	83.0189	58.4906	87.2117	31.0273	80.5031	42.8000
158	82.0755	64.1509	86.3732	33.5430	77.9874	60.4000
178	85.8491	55.6604	87.2117	46.9602	80.5031	57.9000

the data tables. Assume that $\phi(G) = G_b$ on $[-\tau, -2]$ to reflect that the subject has been in fast state and has maintained the glucose level at baseline. Assuming that $\phi(t)$ is linearly increasing from G_b to G_0 in $(-2, 0]$ is in observation that the glucose infusion finishes in 2 minutes and the measurement starts. It is of interest to note that when a rapid increase of secondary insulin secretion is desired, the ramp initial condition produce shorter, impulsive starting values longer time delays; conversely, when a slower increase of secondary insulin secretion must be produced, the ramp starting condition implies longer and impulsive starting values shorter time delays.

In applying the corollaries to these sets of parameter values, we found that the parameters for subject 6 and 8 satisfy both Corollary 4.4 and Corollary 4.5. From the proposed parameter values, we determined numerically that an upper bound for the delay for subject 6 is 8.35226 minutes, and 49.2619 minute for subject 8. However, parameters of subject 7, 13 and 27 do not satisfy either corollary, although the global stability of the equilibrium appears obvious by numerical simulation. We noticed in simulations that conditions of the corollaries are not satisfied due to the rough estimate of the upper bound M_G in Lemma 4.2. We suspect that a better estimate of M_G would improve the scope of applications of Corollary 4.4 and Corollary 4.5.

It is worthwhile to stress the role of the parameter value e . In [14] and [16], the insulin-independent glucose elimination rate parameter e is set to zero. However, from

TABLE 5.2

Experimental data published in [16]. The first column is the time in minute to sample the blood with a two-minute shift. The second and third columns are the data for subject 13. The first valley is at 18' mark. The fifth and sixth columns are the data for subject 27. The first valley is at about 12' mark.

min	G(mg/dl)	I(μ U/l)	min	G(mg/dl)	I(μ U/l)
-2	74.20	24.0	-2	86.47	44.00
0	183.40	231.0	0	345.90	1036.00
2	171.90	127.5	2	275.64	1067.00
4	164.80	124.5	4	263.03	914.00
6	164.10	146.0	6	241.41	415.00
8	150.10	102.5	8	228.80	455.00
10	140.00	129.0	10	227.90	404.00
13	135.10	92.0	12	218.89	216.00
			16	208.98	344.00
18	136.20	88.5	19	199.97	282.00
23	127.00	113.5	22	192.77	232.00
28	118.90	179.5	28	175.65	294.00
			33	163.94	193.00
38	100.90	126.5	38	157.64	227.00
			43	149.53	210.00
48	90.10	91.5	48	147.73	188.00
58	83.40	64.0	58	132.41	116.00
68	79.10	34.0	68	108.99	194.00
78	76.40	30.0	78	97.28	154.00
			88	93.68	95.00
			98	89.18	72.00
118	73.30	28.0	118	84.67	50.00
138	77.30	27.0	138	79.27	38.00
158	76.40	33.5	158	72.06	36.00
178	73.00	23.5	178	72.06	33.00

a physiological point of view, insulin-independent glucose elimination rate could be greater than zero ([5], [18]), so that this term should not simply be ignored.

6. Discussion. The length of delay in a delay differential equation model often plays a critical role in that large delays can destabilize the system. Appropriately determining the range of the delay based on physiology and clinical data is important in theoretical study. In this paper, we suggest that the value of the delay parameter should be set at approximately the time mark of the first clear trough in insulin data. This would help the clinician to assess immediately, even though roughly, this value in applications. Clearly, a full statistical estimation of model parameters based on an appropriate loss functional and an efficient optimization algorithm would provide a more accurate and reliable value for the time delay. Our simulations reveal that the delay may be slight greater than the time to the first clear trough in the insulin concentration time course (refer to the data tables and simulation figures in Section 5). However, even accurate statistical identification of the time delay value from experimental data would not always allow to draw definite conclusions on the stability of the system in a given subject.




FIG. 5.1. Profiles of subject 6 in [4] produced by the model (3.1). The proposed parameters satisfy the condition (4.8) in Corollary 4.4, and both condition (4.11) in Corollary 4.5. The upper bound computed from Corollary 4.4 is 12.2539; while the bound computed from Corollary 4.5 is 15.5167.




FIG. 5.2. Profiles of subject 8 in [4] produced by the model (3.1). The proposed parameters satisfy the condition (4.8) in Corollary 4.4, and both condition (4.11) in Corollary 4.5. The upper bound computed from Corollary 4.4 is 39.5225; while the bound computed from Corollary 4.5 is 49.2619.

Sometimes experimental observations do not allow the investigator to detect a clear first trough in insulinemia for some subjects. For example, there seems no secondary peak or trough for subject #8, but there are seemingly multiple peaks of insulin secretion for subject #27, one at 12' and another one at 22' according to Table 5.2. Also, the actual secondary insulin secretion peak may not be detectable for some subjects: for example, while the simulation for the subject #8 is near-ideal (Fig. 5.2), the profile for subject #27 does not show a peak (Fig. 5.5).

In performing numerical studies for delay differential equation systems, in many cases, particularly for short time dynamics like the IVGTT models, using reasonable functional initial conditions is essential. In this paper, we made use of ramp initial conditions on glycemia, which are more plausible than impulsive jumps to the




FIG. 5.3. Profiles of subject 7 in [4] produced by the model (3.1). The delay parameter τ is out of the upper bounds computed from Corollary 4.4 and Corollary 4.5.




FIG. 5.4. Profiles of subject 13 in [16] produced by the model (3.1). The delay parameter τ is out of the upper bounds computed from Corollary 4.4 and Corollary 4.5.




FIG. 5.5. Profiles of subject 27 in [16] produced by the model (3.1). The delay parameter τ is out of the upper bounds computed from Corollary 4.4 and Corollary 4.5.




FIG. 6.1. Profiles of subject 13 in [4] produced by the model (3.1). The estimated parameters satisfy the condition (4.8) in Corollary 4.4, and both condition (4.11) in Corollary 4.5. The upper bound computed from Corollary 4.4 is 8.93698; while the bound computed from Corollary 4.5 is 17.2833.

highest concentration value adopted until now. This approach should produce more convincing numerical results in clinical applications.

It is important to find out delay dependent condition of the global stability for the model (3.1), under which the clinical data satisfy the conditions. According to Theorem 4.3 and its corollaries, adding a term of insulin-independent glucose uptake, although small, helps considerably in ensuring global stability of the equilibrium, even though AIC does not support this addition from a mere statistical viewpoint, and even though the physiological significance of this term is debatable. Indeed, setting $e = 0$ or $e > 0$ small might not affect the remaining parameter point estimates to any substantial degree. Since a small value of insulin-independent glucose tissue uptake cannot be excluded physiologically according to [5] and [18], we add it to the model in order to exploit the mathematical advantage, which it offers, in satisfying the conditions of both corollaries.

Acknowledgment The authors are grateful to the anonymous reviewers for their invaluable comments that improve the presentation of this manuscript.

REFERENCES

- [1] D.L. BENNETTE AND S.A. GOURLEY, GLOBAL STABILITY IN A MODEL OF THE GLUCOSE/INSULIN INTERACTION WITH TIME DELAY, *Eur. J. Appl. Math.*, 15 (2004) 203.
- [2] R. N. BERGMAN, Y. Z. IDER, C. R. BOWDEN, C. COBELLI, *Quantitative estimation of insulin sensitivity*, *Am. J. Physiol.*, 1979, 236:E667-E677.
- [3] CHERRINGTON, A.D., SINDELAR, D., EDGERTON, D., STEINER, K., MCGUINNESS, O.P., *Physiological consequences of phasic insulin release in the normal animal*. *Diabetes* 51 (2002) (Suppl. 1), S103–S108.
- [4] A. DE GAETANO AND O. ARINO, *Mathematical modeling of the intravenous glucose tolerance test*, *J. Math. Biol.*, 40 (2000), 136–168.
- [5] R. DUHLMEIER, A. HACKER, A. WIDDEL, W. VON ENGELHARDT, AND H.-P. SALLMANN, *Mechanisms of insulin dependent glucose transport into porcine and bovine skeletal muscle*, *Am J. Physiol. Regul. Integr. Comp. Physiol.*, 289: R187–R197, 2005.

- [6] D. V. GIANG, Y. LENBURY, A. DE GAETANO AND P. PALUMBO, *Delay Model of Glucose-Insulin Systems: Global Stability and Oscillated Solutions Conditional on Delays*, JMAA, **343**(2) (2008), pp. 996–1006.
- [7] Y. Kuang, “Delay Differential Equations with Applications in Population Dynamics.” Vol. **191** in the series of Mathematics in Science and Engineering, Academic Press, Boston (1993).
- [8] J. Li and Y. Kuang, *Analysis of a glucose-insulin regulatory models with time delays*. SIAM J. Appl. Math., **67**(3) (2007), 757–776.
- [9] J. Li, Y. Kuang, and B. Li, *Analysis of IVGTT glucose-insulin interaction models with time delay*. Discrete Contin. Dyn. Syst. Ser. B., **1** (2001), 103–124.
- [10] J. Li, Y. Kuang, and C. Mason, *Modeling the glucose-insulin regulatory system and ultradian insulin secretory oscillations with two time delays*. J. Theor. Biol., **242** (2006), 722–735.
- [11] A. MUKHOPADHYAY, A. DE GAETANO AND O. ARINO, *Modelling the intra-venous glucose tolerance test: a global study for a single distributed delay model*, Discr Cont Dyn Syst B, (2004), pp. 407-417.
- [12] A. Makroglou, J. Li, and Y. Kuang, *Mathematical models and software tools for the glucose-insulin regulatory system and diabetes: an overview*. Appl. Numerical Math., **56** (2006), 559–573.
- [13] R. V. OVERGAARD, J. E. HENRIKEN AND H. MADSEN, *Insights to the minimal model of insulin secretion through a mean-field beta cell model*, J. Theor. Biol., **237** (2005), 382-389.
- [14] P. PALUMBO, S. PANUNZI AND A. DE GAETANO, *Qualitative behavior of a family of delay-differential models of the glucose-insulin system*, Discrete and Continuous Dynamical Systems - Series B, Vol.7, No.2, pp.399424, 2007.
- [15] S. PANUNZI, A. DE GAETANO AND G. MINGRONE, *Advantages of the single delay model for the assessment of insulin sensitivity from the intravenous glucose tolerance test*, Theoretical Biology and Medical Modelling, **7**(9), 1–20, 2010
- [16] S. PANUNZI, P. PALUMBO AND A. DE GAETANO, *A discrete Single Delay Model for the Intra-Venous Glucose Tolerance Test*, Theoretical Biology and Medical Modelling, 4:35 (2007), doi:10.1186/1742-4682-4-35.
- [17] N. Pørksen, M. Hollingdal, C. Juhl, P. Butler, J. D. Veldhuis, and O. Schmitz, *Pulsatile insulin secretion: detection, regulation, and role in diabetes*. Diabetes, **51** (2002), S245–S254.
- [18] B. M. SEBASTIAN AND LAURA E. NAGY, *Decreased insulin-dependent glucose transport by chronic ethanol feeding is associated with dysregulation of the Cbl/TC10 pathway in rat adipocytes*, Am J. Physiol Endocrinol Metab 289:1077-1084, 2005. First published Aug 16, 2005; doi:10.1152/ajpendo.00296.2005
- [19] C. SIMON AND G. BRANDENBERGER, *Ultradian oscillations of insulin secretion in humans*, Diabetes. **51** (2002), S258–S261.
- [20] L. F. SHAMPINE AND S. THOMPSON, *Solving DDEs in MATLAB*, Appl. Numer. Math., **37** (2001), pp. 441–458.
- [21] HAL SMITH, *An Introduction to Delay Differential Equations with Applications to the Life Sciences*, Springer, 2010.
- [22] J. Sturis, K. S. Polonsky, E. Mosekilde, and E. Van Cauter, *Computer model for mechanisms underlying ultradian oscillations of insulin and glucose*. Am. J. Physiol., **260** (1991), E801–E809.
- [23] G. TOFFOLO, R. N. BERGMAN, D. T. FINEGOOD, C. R. BOWDEN, C. COBELLI, *Quantitative estimation of beta cell sensitivity to glucose in the intact organism: a minimal model of insulin kinetics in the dog*, Diabetes 1980, 29:979-990.
- [24] I. M. Tolic, E. Mosekilde and J. Sturis, *Modeling the insulin-glucose feedback system: The significance of pulsatile insulin secretion*. J. Theor. Biol., **207** (2000), 361–375.
- [25] B. Topp, K. Promislow, G. De Vries, R. M. Miura, and D. T. Finegood, *A Model of β -cell mass, insulin, and glucose kinetics: pathways to diabetes*, J. Theor. Biol., **206** (2000), 605–619.
- [26] H. Wang, J. Li, and Y. Kuang, *Mathematical modeling and qualitative analysis of insulin therapies*, Math. Biosci., **210** (2007), 17–33.
- [27] H. Wang, J. Li, and Y. Kuang, *Enhanced modeling of the glucose-insulin system and its applications in insulin therapies*, J. Biological Dynamics, **3** (2009), 22-38.